HEALTH BASICS

How much sleep you need

Sleep need is the amount of regular sleep that supports health and daytime alertness.

WHAT IT IS
The sleep time your body needs for health and alertness

WHERE IT COMES UP
Daily routines, parenting, shift work, and sleep concerns

WHY IT MATTERS
Supports attention, mood, metabolism, immunity, and physical safety

WHAT IT IS NOT
Not a disease or diagnosis

WHO TO ASK
Primary care clinician or sleep medicine specialist

HOW IT IS CHECKED
Sleep diary, daytime function, and sometimes a sleep study

Overview

Sleep need is the amount of sleep that helps your brain and body function well. It is not a disease or a diagnosis.

Age is the main guide to how much sleep people generally need. Your needs may also be affected by illness, pregnancy, physical activity, sleep loss, and the quality and timing of your sleep.

Recommendations describe a healthy range for most people, not an exact number that everyone must reach. The American Academy of Sleep Medicine (AASM) recommends the following amounts:

Sleep in babies younger than 4 months is often spread across the day and night. Their patterns change quickly, so these consensus recommendations do not give one target for this age group. A pediatric clinician can answer questions about an individual baby's sleep and feeding pattern.

- Ages 4 to 12 months: 12 to 16 hours per 24 hours, including naps
- Ages 1 to 2 years: 11 to 14 hours, including naps
- Ages 3 to 5 years: 10 to 13 hours, including naps

- Ages 6 to 12 years: 9 to 12 hours
- Ages 13 to 18 years: 8 to 10 hours
- Adults ages 18 to 60: at least 7 hours each night
- Adults ages 61 to 64: 7 to 9 hours
- Adults age 65 and older: 7 to 8 hours

How it works

Sleep is an active process. During the night, your brain moves through several stages, including rapid eye movement (REM) sleep and non-REM sleep. These stages support learning, memory, emotional processing, tissue repair, immune function, and hormone control.

Sleep duration means time actually spent asleep. It is not always the same as time in bed. For example, eight hours in bed may include time spent reading, trying to fall asleep, or lying awake during the night.

Timing also matters. Your circadian rhythm is the internal body clock that helps control when you feel awake or sleepy. Light, work hours, meals, activity, and travel can shift this rhythm. A person may get enough hours but still feel poorly rested if sleep happens at changing times or is often interrupted.

Sleep quality matters as well. Loud snoring, breathing pauses, pain, hot flushes, caregiving duties, or an uncomfortable environment can repeatedly disturb sleep. A sleep tracker may estimate movement or sleep stages, but consumer devices cannot measure sleep need precisely or diagnose a sleep disorder.

You may need more sleep for a time after missing sleep, during recovery from illness, or during periods of greater physical demand. Regularly needing much more sleep than expected is worth discussing with a clinician, especially if you remain sleepy during the day.

Why it matters for your health

Adequate sleep supports attention, reaction time, judgment, mood, and coordination. It also plays a role in blood pressure, blood sugar control, appetite, immune responses, and growth in children and teenagers.

Repeatedly getting too little sleep can make learning, work, and driving more difficult. It is associated with a higher risk of injuries and several long-term health conditions, including high blood pressure, heart disease, type 2 diabetes, depression, and obesity. These associations do

not mean that sleep alone causes every condition.

Children and teenagers may show insufficient sleep through difficulty waking, falling asleep in class, irritability, or problems concentrating. These changes can have many causes, so they do not confirm that sleep loss is the reason.

How you function during the day provides useful context. Feeling alert through ordinary activities, waking without extreme difficulty, and not needing frequent unplanned naps may suggest that your sleep amount and quality are adequate. However, people can become used to feeling tired, so daytime function is not a perfect test.

What can go wrong

Not allowing enough time for sleep can lead to sleep deprivation. This may happen because of work, school, caregiving, social schedules, screen use, or repeated nighttime interruptions.

Sometimes enough time is available, but another condition affects sleep. Examples include:

Medicines, alcohol, caffeine, pain, anxiety, depression, menopause, pregnancy, and some medical conditions can also change sleep or daytime alertness. Sleeping longer than usual does not by itself prove that something is wrong. It can reflect recovery, a changed schedule, medication effects, or illness.

Arrange routine medical follow-up if sleep problems continue, you regularly need much more sleep than before, or tiredness interferes with daily life. Also seek advice for loud habitual snoring, witnessed breathing pauses, waking with choking, or repeated unintended sleep episodes.

Do not drive or operate machinery if you are struggling to stay awake. Stop somewhere safe and arrange another way to travel. Call emergency services for severe breathing difficulty, blue or grey lips, new confusion, collapse, or if someone cannot be awakened normally.

- Insomnia: ongoing difficulty falling asleep, staying asleep, or returning to sleep
- Obstructive sleep apnea: repeated narrowing or closure of the airway during sleep
- Restless legs syndrome: an urge to move the legs, often with uncomfortable sensations
- Narcolepsy: a neurological condition that affects control of sleep and wakefulness
- Circadian rhythm sleep-wake disorders: a mismatch between the body clock and the required schedule

What you can do

There is no blood test or single calculation that reveals your exact sleep need. A useful first step is to compare age-based recommendations with your schedule and daytime function.

Keep a sleep diary for one or two weeks. Record:

Try to allow enough time for the recommended amount of sleep, not just that amount of time in bed. A fairly consistent wake time can help your body clock. If work, caregiving, housing, pain, or another practical issue limits sleep, focus on changes that are realistic rather than treating the target as a personal failure.

Ask a clinician whether a medicine or health condition could be affecting sleep if the pattern is new or persistent. Bring your sleep diary and, if relevant, observations from someone who has noticed snoring or breathing pauses.

A sleep study may be used when a sleep disorder is suspected. It records information such as breathing, oxygen level, heart rate, movement, and brain activity. It does not simply calculate how many hours you personally need.

- When you went to bed and when you tried to sleep
- Estimated time asleep and awake during the night
- Wake time, naps, and alarm use
- Caffeine, alcohol, shift work, and medicines that may affect sleep
- Daytime sleepiness and any unintended dozing

Common questions

Can some adults be healthy with less than seven hours?

Individual sleep need differs, and a small number of people naturally sleep less. However, regularly sleeping under seven hours is not recommended for most adults. Feeling accustomed to short sleep does not always mean attention and reaction time are unaffected.

Is nine hours too much for an adult?

Not necessarily. More than nine hours may be appropriate for young adults, people recovering from sleep loss, and people who are ill. A clear increase in sleep time, persistent daytime sleepiness, or difficulty functioning deserves routine medical review.

Can naps replace nighttime sleep?

Naps count toward total sleep for young children. For adults, a short nap may improve alertness, but it may not fully replace regular nighttime sleep. Long or late naps can also make

it harder to fall asleep at night.

Should I trust my sleep tracker?

Use it to notice broad patterns rather than exact sleep stages or diagnoses. Consumer trackers estimate sleep from signals such as movement and heart rate. If the device's result conflicts with how you feel, discuss the overall pattern with a clinician rather than relying on one score.

Can I catch up on sleep at weekends?

Extra sleep may reduce some immediate sleepiness after a short week. It does not reliably remove all effects of ongoing sleep loss, and large changes in wake time can shift your body clock. The more useful goal is enough sleep on a regular basis.

Sources

- American Academy of Sleep Medicine: Recommended amount of sleep for pediatric populations , Paruthi S and colleagues, Journal of Clinical Sleep Medicine , 2016
- American Academy of Sleep Medicine and Sleep Research Society: Recommended amount of sleep for a healthy adult , Watson NF and colleagues, Journal of Clinical Sleep Medicine , 2015
- Centers for Disease Control and Prevention: About Sleep
- National Heart, Lung, and Blood Institute: How Much Sleep Is Enough?
- American Academy of Sleep Medicine: Drowsy Driving Health Advisory
- American Academy of Sleep Medicine: Clinical practice guideline for diagnostic testing for adult obstructive sleep apnea , Kapur VK and colleagues, Journal of Clinical Sleep Medicine , 2017