HEALTH BASICS

Interaction checker

An interaction checker identifies possible problems between medicines, supplements, foods, and health conditions.

WHAT IT IS
A tool that screens for possible medicine interactions

WHERE IT COMES UP
Starting, stopping, or combining medicines, foods, and supplements

WHY IT MATTERS
Helps identify combinations needing avoidance, adjustment, or monitoring

WHAT IT IS NOT
Not a diagnosis or guarantee of medicine safety

WHO TO ASK
A pharmacist or prescribing clinician

HOW IT IS CHECKED
Search of current medicines in a trusted interaction database

Overview

An interaction checker is a reference tool. It compares medicines and other substances to identify combinations that may change how they work or increase the chance of harm.

It is not a disease, diagnosis, or medical test. It also cannot prove that an interaction will happen. A warning means that the combination deserves closer review.

Depending on the checker, it may look for interactions involving:

Checkers are used by pharmacists, prescribers, hospitals, and members of the public. Professional systems may include more detailed information than free consumer tools.

An interaction result should be interpreted in context. Your doses, age, kidney and liver function, pregnancy status, other health conditions, and reason for treatment can all affect what a warning means.

- Prescription and over-the-counter medicines
- Vitamins, minerals, and herbal products
- Alcohol, cannabis, or certain foods

- Health conditions that may make a medicine less suitable

How it works

An interaction checker searches a database for known or suspected conflicts. Information may come from clinical studies, official medicine labels, case reports, and monitoring after a medicine reaches the market.

Some interactions affect what the body does to a medicine. For example, one substance may reduce absorption from the gut or slow the liver's breakdown of another medicine. This can make the medicine level too low or too high.

Other interactions affect what medicines do to the body. Two medicines may both cause drowsiness, lower blood pressure, increase bleeding, or slow breathing. Taking them together can strengthen that shared effect.

A checker commonly reports the possible effect, the evidence behind it, and a severity category. It may suggest avoiding the combination, separating doses, changing a dose, or arranging monitoring. These are prompts for review, not personal treatment instructions.

Severity labels are not standardized. One checker may call an interaction "major," while another calls it "moderate." Databases may also differ in how quickly they add new evidence or whether they include foods, supplements, and health conditions.

Many checkers cannot fully account for:

Correct entry matters. Brand names may contain more than one active ingredient, and similar names can refer to different medicines. The route also matters: a swallowed medicine may behave differently from a cream, injection, or inhaled product.

- The exact dose, formulation, or timing
- A recently changed medicine
- Kidney or liver function
- Previous reactions or individual sensitivity
- Several medicines acting together
- Limited evidence about supplements

Why it matters for your health

Interactions can change whether treatment works and how likely it is to cause unwanted

effects. This becomes more important as the number of medicines and supplements increases, because there are more possible combinations to review.

A checker can help you notice an issue before starting a new product. It can also give you a more specific question to ask, such as whether doses should be separated or whether a blood test is needed.

Not every listed interaction requires a change. Some combinations are used deliberately because their benefits outweigh the risks. They may be reasonable with a particular dose, clear instructions, or routine monitoring.

The reverse is also important: a checker showing "no interaction found" does not prove that a combination is safe. The interaction may be uncommon, newly recognized, missing from that database, or related to a health condition the tool did not consider.

What can go wrong

A checker can miss an interaction or produce so many alerts that an important one is overlooked. Results may be misleading if your medicine list is incomplete or a combination product is entered incorrectly.

Possible effects of specific medicine combinations include bleeding, very low blood pressure, low blood sugar, abnormal heart rhythms, excessive sedation, slowed breathing, kidney injury, or liver injury. Some combinations can also reduce treatment effectiveness.

Certain interactions relate to named medical emergencies. For example, some medicine combinations can contribute to serotonin syndrome, a serious reaction involving excess serotonin activity, or severe central nervous system depression, in which breathing and alertness become dangerously reduced. These outcomes are linked to particular combinations; they are not effects of the checker itself.

Do not stop a prescribed medicine solely because a website displays a warning. Suddenly stopping some medicines can cause withdrawal effects or allow the condition being treated to worsen.

Routine follow-up is appropriate when you find a warning but feel well. A pharmacist or prescribing clinician can check whether it applies to your dose and circumstances.

Seek urgent medical advice for a new concerning effect after starting or combining medicines, such as unusual bleeding, repeated fainting, marked confusion, severe restlessness with fever, or a very fast or irregular heartbeat.

Call emergency services for trouble breathing, collapse, a seizure, severe chest pain, signs of a

stroke, or if someone cannot be awakened. Contact a local poison control service promptly after a suspected overdose or accidental medicine combination, even if there are no symptoms yet.

What you can do

Keep one current medicine list. Include prescriptions, over-the-counter products, vitamins, herbal products, and medicines used only occasionally. Record the active ingredient, strength, dose, route, and usual schedule when known.

Bring the list to appointments and pharmacy visits. Update it when anything starts, stops, or changes. After a hospital stay, compare your new instructions with your previous list because unplanned duplication can occur during transitions in care.

Before adding a medicine or supplement, you can ask a pharmacist:

Using one pharmacy when practical can make it easier for the pharmacist's system to screen your full prescription record. It will not capture products bought elsewhere unless you report them.

There is no single blood test that detects all interactions. When a particular interaction is relevant, monitoring might include blood pressure, blood sugar, an electrocardiogram that records heart rhythm, blood tests for kidney or liver function, or measurement of a medicine level. Many combinations need no testing.

If you use an online checker, enter each active ingredient rather than relying only on brand names. Check whether the tool includes supplements, foods, alcohol, and health conditions. Treat the result as a starting point for review.

Record any new effect, when it began, and how it relates to your doses. Do not test a suspected interaction by taking the combination again.

- Does this interact with anything on my complete list?
- Does the warning apply to my dose and how I use it?
- Should I avoid the combination or separate the doses?
- Do I need symptom monitoring, a blood test, or another check?
- Which effects would require urgent help?

Common questions

Does a "major" warning mean I must stop the medicine?

Not necessarily. It means the combination may have an important risk. A pharmacist or prescribing clinician should review why each medicine is being used, what alternatives exist, and whether monitoring can reduce the risk. Unless you have emergency symptoms, seek advice before making a sudden change.

Why do two interaction checkers give different answers?

They may use different evidence, severity categories, update schedules, or rules. One may include theoretical interactions that another excludes. When results conflict, ask a pharmacist to check professional references and the official prescribing information.

Can food interact with medicine?

Yes. Certain foods can alter absorption or metabolism, while a medicine may need to be taken with or without food. Grapefruit and foods containing vitamin K are relevant to some medicines, but not all. Follow the instructions for your specific product rather than avoiding foods unnecessarily.

Do herbal products and vitamins count?

Yes. Supplements can change medicine levels or add to effects such as bleeding or drowsiness. Evidence is often less complete than it is for prescription medicines, so an empty result does not confirm safety.

Can I check medicines for another person?

You can screen the list, but the result may not reflect that person's health conditions, doses, or laboratory results. Their pharmacist or prescribing clinician should interpret important warnings.

Is an interaction checker useful for one medicine?

It can be if the checker includes food, alcohol, supplements, or health-condition warnings. A single medicine can also contain several active ingredients, especially cold, pain, and sleep products.

Sources

- US Food and Drug Administration. Drug Interactions: What You Should Know .
- US Food and Drug Administration. Clinical Drug Interaction Studies-Cytochrome P450 Enzyme- and Transporter-Mediated Drug Interactions: Guidance for Industry . 2020.
- National Library of Medicine. DailyMed: About DailyMed .

- National Center for Complementary and Integrative Health. Herb-Drug Interactions .
- Kheshti R, Aalipour M, Namazi S. A comparison of five common drug-drug interaction software programs regarding accuracy and comprehensiveness . Journal of Research in Pharmacy Practice. 2016;5(4):257-263.
- [Roblek T, Vaupotic T, Mrhar A, Lainscak M. Drug-drug interaction software in clinical practice: a systematic review . European Journal of Clinical Pharmacology. 2015;71:131-142.](https://doi.org/10.1007/s00228-014-178 softwares)