HEALTH BASICS

Prosthetic graft

A prosthetic graft is a manufactured tube or patch used to carry blood or repair a blood vessel.

WHAT IT IS
A manufactured tube or patch replacing or bypassing tissue

WHERE IT COMES UP
Vascular surgery, dialysis access, imaging, or operation reports

WHY IT MATTERS
Restores blood flow but may require long-term monitoring

WHAT IT IS NOT
Not a disease or diagnosis

WHO TO ASK
Vascular surgeon, cardiac surgeon, or dialysis access team

HOW IT IS CHECKED
Examination, duplex ultrasound, CT, or location-specific imaging

Overview

A prosthetic graft is a manufactured medical implant. Surgeons most often use one to replace, repair, or bypass part of a blood vessel. This article focuses on vascular prosthetic grafts.

A graft may reroute blood around a narrowed or blocked artery. It may replace part of an enlarged or damaged aorta, the main artery leaving the heart. A graft can also connect an artery to a vein to create access for hemodialysis, a treatment that filters the blood when the kidneys cannot.

Common graft materials include polyester, sometimes called Dacron, and expanded polytetrafluoroethylene, or ePTFE. These materials are designed to stay in the body. The shape may be a straight tube, a branching tube, or a patch.

"Prosthetic" means manufactured. It does not mean an artificial limb. A prosthetic graft also differs from a graft made from your own vein or from donated human tissue.

A stent graft is related but not identical. It combines graft fabric with a metal framework and is usually placed through a blood vessel using a catheter. Reports may use terms such as vascular graft , synthetic graft , bypass graft , or endograft , so it is useful to ask exactly which device you have.

How it works

For an open vascular operation, the surgeon usually attaches each end of the graft to a blood vessel with stitches. These joins are called anastomoses. Blood then travels through the graft instead of through the damaged, enlarged, or blocked section.

A bypass graft does not remove the underlying artery disease. It creates another route for blood flow. An aortic replacement graft takes the place of a section of the aorta. A patch widens or repairs part of a vessel rather than replacing it with a full tube.

A dialysis graft is placed under the skin, usually in an arm. It connects an artery to a vein and provides an area where dialysis needles can be inserted after the access has healed and the dialysis team says it is ready.

Blood is naturally more likely to clot when it contacts an artificial surface than when it flows through a healthy blood vessel. Graft size, location, blood flow, and the condition of nearby vessels all affect how well a graft functions.

Some people are prescribed an antiplatelet medicine or an anticoagulant like warfarin or apixaban , sometimes called a blood thinner. The right medicine is based on the type of operation, graft location, bleeding risk, and other health conditions. Do not start, stop, or change one because of general information about grafts.

Why it matters for your health

A prosthetic graft can restore or protect blood flow when a person's own vessel is unsuitable for repair. The health effect therefore comes from where the graft is placed and why it was needed.

Follow-up is not the same for every graft. A graft in the leg may be checked for changes in circulation. An aortic graft may need imaging to assess the repaired aorta and nearby blood vessels. A dialysis graft is checked during dialysis and before needles are inserted.

Checks may include feeling pulses, examining the skin, measuring blood pressure in different parts of the body, or listening to blood flow. Duplex ultrasound uses sound waves to show blood movement and possible narrowing. CT or MRI may be used for deeper grafts, particularly in the chest, abdomen, or pelvis.

Some grafts need scheduled surveillance even when you feel well. Other follow-up is guided by the operation and your overall vascular health. Your discharge letter, implant record, or surgical team should state what applies to you.

What can go wrong

A prosthetic graft is not an illness, but several medical problems can affect it.

Narrowing or blockage: Scar tissue or continuing blood vessel disease can reduce blood flow near a graft. A clot may partly or completely block it. This is called graft thrombosis.

Infection: Bacteria can infect the graft or tissue around it. Infection may appear soon after surgery or later. Possible signs include increasing redness, warmth, swelling, pain, drainage, an opening in the wound, or fever. These signs have other possible causes, but they need prompt assessment.

Problems at a join: An anastomosis may narrow, leak, or weaken. A contained leak can form a pulsing swelling called a pseudoaneurysm, meaning blood has collected outside the vessel wall but remains connected to the bloodstream.

Bleeding or rupture: Serious bleeding is uncommon but can occur from a damaged graft, a weakened join, or repeated needle sites in a dialysis graft.

Fluid around the graft: Clear fluid can sometimes collect near a graft. This is called a seroma. A new or growing swelling should be examined rather than assumed to be harmless.

Dialysis access problems: A dialysis graft may clot, become infected, develop a narrowed area, or send too much blood away from the hand. Reduced hand circulation is sometimes called steal syndrome. A cold, painful, weak, or numb hand needs urgent assessment.

A connection between an aortic graft and the bowel, called an aortoenteric fistula, is a rare but dangerous cause of internal bleeding or infection.

Call emergency services for:

Apply firm direct pressure to external bleeding while emergency help is being arranged. Do not press on or attempt to push back a new pulsing lump.

Seek urgent same-day medical advice for fever with redness, drainage, worsening pain, or swelling over a graft. A new lump, a change in limb circulation, or a missing dialysis graft vibration also needs prompt assessment.

- Sudden severe pain, paleness, coldness, numbness, or weakness in a limb
- Heavy or uncontrolled bleeding from a graft or dialysis needle site
- Sudden severe chest, back, or abdominal pain, especially with fainting
- Vomiting blood, passing black stools, or collapsing

What you can do

Keep a record of the graft's location, purpose, operation date, and device details if available. Carry an implant card if you were given one. Share this information before surgery, imaging, or other procedures.

Ask your surgical or dialysis team:

Take prescribed medicines as directed. Do not stop antiplatelet or anticoagulant treatment before a procedure unless the clinician managing that medicine gives you a plan.

There is no reliable home test for most internal grafts. Check the wound only as instructed during recovery and attend planned follow-up even if you feel well.

If you have a dialysis graft, the access team may teach you to feel its vibration, known as a thrill. Report a clear change or loss of that vibration promptly. Avoid blood pressure cuffs, blood draws, tight clothing, and prolonged pressure on the access arm if your dialysis team has told you to protect it this way.

- Whether routine examination or imaging is planned
- Which changes should lead to a same-day call
- Whether any activity or lifting limits still apply
- Which medicines relate to the graft
- Whom to contact if you move or miss follow-up

Common questions

Can I feel a prosthetic graft?

Most deep grafts cannot be felt. A dialysis graft lies closer to the skin, and you may feel a steady vibration from blood flow. A new tender, enlarging, or pulsing lump is not a normal way to check a graft and should be assessed.

Does a prosthetic graft last for life?

There is no single lifespan for all grafts. Durability is influenced by the graft's material and location, blood flow, nearby vessel disease, infection, and the reason it was placed. Some continue to work for many years, but long-term grafts may still need monitoring.

Can I have an MRI?

Many vascular graft materials can be scanned safely, but some graft systems include metal stents, clips, or other components with specific MRI conditions. Tell the imaging service about the implant and provide the device card or operation details. The imaging team should confirm

compatibility before the scan.

Will I always need a blood thinner?

Not necessarily. Medicine choices differ between aortic repairs, leg bypasses, dialysis grafts, and other procedures. They also reflect your bleeding risk and other conditions. Follow the plan written for you rather than assuming the graft requires lifelong treatment.

Can a blocked graft be treated?

Sometimes. Treatment may involve removing or dissolving a clot, widening a narrowed section with a balloon, inserting a stent, revising the graft, or creating another bypass. The suitable option is determined after assessment and imaging.

Do I need antibiotics like amoxicillin or doxycycline before dental work?

A prosthetic vascular graft does not automatically mean that antibiotics are needed before every dental procedure. Recommendations are based on the implant, timing, procedure, and other medical factors. Ask the clinician responsible for the graft and your dentist rather than taking leftover antibiotics.

Sources

- Zierler RE, Jordan WD, Lal BK, et al. "The Society for Vascular Surgery Practice Guidelines on Follow-up After Vascular Surgery Arterial Procedures." Journal of Vascular Surgery . 2018;68(1):256-284.
- Chakfé N, Diener H, Lejay A, et al. "Editor's Choice - European Society for Vascular Surgery 2020 Clinical Practice Guidelines on the Management of Vascular Graft and Endograft Infections." European Journal of Vascular and Endovascular Surgery . 2020;59(3):339-384.
- Isselbacher EM, Preventza O, Hamilton Black J III, et al. "2022 ACC/AHA Guideline for the Diagnosis and Management of Aortic Disease." Circulation . 2022;146(24):e334-e482.
- Lok CE, Huber TS, Lee T, et al. "KDOQI Clinical Practice Guideline for Vascular Access: 2019 Update." American Journal of Kidney Diseases . 2020;75(4 Suppl 2):S1-S164.
- American College of Radiology. ACR Manual on MR Safety . 2024 edition. Guidance on screening and assessing implanted medical devices before MRI.