HEALTH BASICS

# RICE

RICE is a first-aid approach using rest, ice, compression, and elevation after some minor injuries.

**WHAT IT IS**
A first-aid approach for recent minor soft-tissue injuries

**WHERE IT COMES UP**
Sprains, strains, bruises, and sports injuries

**WHY IT MATTERS**
May ease pain and swelling during early recovery

**WHAT IT IS NOT**
Not a disease or diagnosis

**WHO TO ASK**
Pharmacist, primary care clinician, physiotherapist, or urgent care service

**HOW IT IS CHECKED**
No test; the injury itself may need assessment

## Overview

RICE stands for rest, ice, compression, and elevation . It is a traditional first-aid approach for some recent injuries, especially minor sprains, strains, and bruises.

A sprain affects a ligament, which is a strong band connecting one bone to another. A strain affects a muscle or tendon. Tendons connect muscles to bones. These are known as soft-tissue injuries.

RICE is not a disease, diagnosis, or treatment for every injury. It cannot tell you whether a bone is broken or a tendon is torn. Its main role is to help manage discomfort and swelling while you monitor the injury or arrange an assessment.

Current advice often favors relative rest rather than keeping the area completely still. Gentle movement and a gradual return to activity may support recovery once they are comfortable and appropriate for the injury.

## How it works

Each part of RICE has a different purpose.

Ice appears most useful for short-term pain relief. Research has not clearly shown that it speeds tissue healing. Prolonged or excessive cooling may interfere with normal biological processes involved in repair, so more ice is not necessarily better.

Complete rest can also have drawbacks. Keeping a joint or muscle inactive for too long can contribute to stiffness, weakness, and reduced balance. Depending on the injury, gentle movement followed by gradually increased activity may be more helpful.

You may also see newer terms such as PRICE , POLICE , or PEACE and LOVE . These approaches add ideas such as protection, appropriate loading, education, and exercise. They reflect a shift away from prolonged rest and toward recovery guided by pain, function, and professional advice.

- Rest: Stop or reduce activities that clearly increase pain. This does not always mean complete bed rest or avoiding all movement. A brace, splint, or walking aid may sometimes protect an injury, but it should fit properly and suit the injury.
- Ice: A cold pack can temporarily numb pain and may limit swelling. Wrap it in a damp towel or other fabric rather than putting it directly on your skin. Follow the cold-pack instructions and check your skin regularly.
- Compression: An elastic bandage may help control swelling and provide light support. It should feel snug, not tight. Remove or loosen it if pain increases or the area beyond the bandage becomes numb, cold, pale, or blue.
- Elevation: Raising the injured area above heart level, when practical, may help fluid drain away and reduce swelling.

## Why it matters for your health

RICE gives you a simple way to respond immediately to a minor injury. It may make the first hours or days more comfortable while swelling settles.

However, the acronym can make different injuries seem more alike than they are. A mild ankle sprain, a fracture, and an Achilles tendon rupture can all follow a sudden ankle injury, but they need different care. Using RICE should not delay an assessment when the cause or severity is uncertain.

How quickly you should move the area also differs by injury. Early, controlled movement may help many uncomplicated sprains and strains, while a fracture or major tendon injury may need protection or immobilization. A clinician can advise you if pain, swelling, or loss of function makes the next step unclear.

## What can go wrong

The main concern is mistaking a more serious injury for a minor sprain or strain. Related conditions include fractures, dislocations, major ligament tears, tendon ruptures, deep wounds, and significant muscle tears.

Compression can cause problems if a bandage is too tight. Warning signs include increasing pain, tingling, numbness, or fingers or toes becoming cold, pale, or blue. Remove the bandage promptly if these occur.

Ice can injure the skin if it is applied directly or left in place too long. People with reduced skin sensation or poor circulation may not notice cold injury developing. Ask a clinician whether cold treatment is suitable if either applies to you.

Seek emergency help for:

Seek urgent assessment if you heard a crack, cannot use the limb or put weight on it, have marked swelling or bruising, or develop persistent numbness. A sudden pop with loss of strength or movement may indicate a tendon or ligament injury.

Arrange routine follow-up if pain, swelling, movement, or function is not steadily improving, or if the joint repeatedly gives way. Persistent problems may need examination, rehabilitation, or imaging such as an X-ray.

- A limb that looks severely deformed or has bone visible through a wound
- Heavy bleeding that does not stop with firm pressure
- Skin beyond the injury that remains cold, pale, or blue
- Severe, rapidly worsening pain with a tense or very swollen limb
- Loss of consciousness or a serious head, neck, or back injury

## What you can do

After a minor injury, stop the activity and check what happened. Look for deformity, an open wound, unusual skin color, numbness, or loss of function before assuming RICE is suitable.

If there are no urgent warning signs, you can:

Do not force a painful joint through movement. As discomfort settles, gentle movement and a gradual return to normal activity may be reasonable. The right pace depends on the injured structure and how well it functions, not just how many days have passed.

Ask a pharmacist or clinician how to use a bandage if you are unsure. You can also ask whether

you need an examination, an X-ray, or referral to a physiotherapist. An X-ray shows bones well but does not show every ligament, tendon, or muscle injury.

If you use pain medicine, follow the label and ask a pharmacist whether it is suitable with your health conditions and other medicines.

- Protect the area from movements that sharply increase pain
- Use a wrapped cold pack for brief periods, following its instructions
- Apply a properly fitted elastic bandage if compression is comfortable
- Elevate the area when practical
- Record whether swelling, movement, and ability to use the area improve

## Common questions

Does RICE make an injury heal faster?

That has not been clearly established. Ice and elevation may reduce discomfort or swelling, but evidence that the complete RICE approach speeds healing is limited. Appropriate movement and rehabilitation may be more important for longer-term recovery.

Should I rest until there is no pain?

Not necessarily. Avoid activities that cause sharp or increasing pain, but complete rest for an extended period may lead to stiffness and weakness. A clinician or physiotherapist can help set a safe level of activity after a more significant injury.

Can I put ice directly on my skin?

No. Place fabric between your skin and the cold pack. Stop if the skin becomes very painful, blotchy, unusually pale, or numb.

Is swelling always harmful?

No. Swelling is part of the body's response to injury, although too much swelling can increase pain and limit movement. The goal is usually comfort and useful function, not eliminating every sign of swelling.

Is RICE still recommended?

It remains widely recognized as basic first aid, but it is no longer viewed as a complete recovery plan. Newer approaches emphasize protection when needed, education, gradual loading, exercise, and avoiding unnecessary prolonged rest.

## Sources

- NHS. "Sprains and strains."
- American Academy of Orthopaedic Surgeons. "Sprained Ankle." OrthoInfo.
- van den Bekerom MPJ, et al. "What Is the Evidence for Rest, Ice, Compression, and Elevation Therapy in the Treatment of Ankle Sprains in Adults?" Journal of Athletic Training . 2012;47(4):435-443.
- Kaminski TW, et al. "National Athletic Trainers' Association Position Statement: Conservative Management and Prevention of Ankle Sprains in Athletes." Journal of Athletic Training . 2013;48(4):528-545.
- Dubois B, Esculier JF. "Soft-tissue injuries simply need PEACE and LOVE." British Journal of Sports Medicine . 2020;54:72-73.