HEALTH BASICS

S-adenosylmethinoine

S-adenosylmethionine is a natural body chemical also sold as the supplement SAM-e.

WHAT IT IS
A natural compound involved in essential cell reactions

WHERE IT COMES UP
Supplement labels and discussions about mood, joints, or liver health

WHY IT MATTERS
Helps transfer chemical groups needed for normal cell function

WHAT IT IS NOT
Not a disease or diagnosis

WHO TO ASK
A pharmacist or primary care clinician

HOW IT IS CHECKED
No routine clinical test

Overview

The standard spelling is S-adenosylmethionine , although the name is sometimes misspelled as S-adenosylmethinoine. It is usually shortened to SAM-e , pronounced "sammy."

SAM-e is a compound that your body makes from methionine, an amino acid found in protein, and adenosine triphosphate, or ATP, which supplies energy to cells. It is present throughout the body and is especially important in the liver and brain.

This compound is not a disease, diagnosis, vitamin, hormone, or prescription antidepressant like sertraline or fluoxetine . It is also different from methionine, homocysteine, and serotonin, although it takes part in chemical pathways connected with these substances.

Manufactured SAM-e is sold as a dietary supplement in the United States. In some other countries, certain forms are regulated as medicines. Product standards and approved uses therefore differ by country. In the United States, dietary supplements are not approved by the Food and Drug Administration for effectiveness before they are marketed.[1,2]

SAM-e supplements are promoted for depression, osteoarthritis, and liver conditions. Research has not established that an over-the-counter supplement reliably treats these conditions. It should not replace an assessment or treatment that has been recommended for a diagnosed health problem.[1,3,4]

How it works

SAM-e mainly acts as a methyl donor . A methyl group is a small collection of carbon and hydrogen atoms. Moving methyl groups between molecules is called methylation.

Cells use methylation to help:

After SAM-e donates a methyl group, it becomes S-adenosylhomocysteine and then homocysteine. Homocysteine can be recycled into methionine or directed into pathways that help make glutathione, an antioxidant used by cells.[5,6]

Folate, vitamin B12, and vitamin B6 help parts of this wider cycle work. However, taking SAM-e is not a substitute for identifying or treating a deficiency in one of these vitamins.

The body tightly coordinates these reactions. A higher supplement intake does not simply cause every methylation process to work better. Oral SAM-e must also survive digestion and be absorbed before it can enter the body's chemical pathways.

- Regulate how genes are switched on and off
- Make or process certain brain-signalling chemicals
- Build parts of cell membranes
- Process proteins, fats, medicines, and other substances

Why it matters for your health

Normal SAM-e production is part of basic cell function. Researchers are interested in it because methylation and related pathways affect the nervous system, joints, liver, and many other tissues.

For depression, some studies suggest that SAM-e may have an antidepressant effect, either alone or when added to prescribed treatment. However, trials have generally been small or inconsistent. A Cochrane review found that the available evidence was not strong enough to support firm conclusions about benefit or safety.[3]

For osteoarthritis, older studies examined whether SAM-e could reduce pain or improve joint function. The evidence has limitations, and it does not show clearly that SAM-e provides dependable benefit comparable with established treatments.[1,4]

SAM-e has also been studied in several liver and bile-flow conditions. These are distinct medical problems with different causes. Research into a body pathway does not mean that a shop-bought supplement can treat every condition involving that pathway.[1,5]

Supplement labels may use terms such as "mood support," "joint health," or "liver support." These are broad marketing descriptions. They are not proof that a product has been shown to treat depression, arthritis, or liver disease.

What can go wrong

SAM-e supplements can cause unwanted effects. Reported effects include nausea, diarrhoea, constipation, stomach discomfort, headache, dizziness, restlessness, anxiety, and difficulty sleeping.[1] The amount of active ingredient and the way tablets release it may differ between products.

SAM-e may increase serotonin activity. Combining it with antidepressants or other substances that affect serotonin could raise the risk of serotonin syndrome , a potentially serious reaction. Relevant medicines may include some antidepressants and certain medicines used for migraine, pain, or cough. Supplements such as St John's wort may also affect serotonin.

Seek emergency help if, after taking a new medicine or supplement, you develop a combination of marked agitation or confusion, fever, heavy sweating, diarrhoea, shaking, severe muscle stiffness, or loss of coordination. These symptoms can have several causes and need prompt assessment.

SAM-e may trigger mania or hypomania in some people with bipolar disorder. These are periods of unusually elevated or irritable mood, increased activity, reduced need for sleep, rapid speech, or impulsive behaviour. Contact a mental health service promptly if these changes appear. Severe agitation, dangerous behaviour, psychosis, or thoughts of suicide require urgent or emergency support.

Using SAM-e to manage persistent low mood can also delay an assessment for depression, bipolar disorder, medicine effects, thyroid problems, or other causes. Routine follow-up is appropriate for ongoing low mood without immediate danger. Thoughts of harming yourself or being unable to stay safe require immediate crisis or emergency help.

Safety information during pregnancy and breastfeeding is limited. Product quality may also be uncertain, particularly when supplements are bought from sellers without clear manufacturing or independent testing information.[1,2]

What you can do

You do not need to measure or increase SAM-e as part of routine health care. There is no standard clinical blood test used to check whether most people have a "SAM-e deficiency."

If you are considering a supplement, bring the exact product label to a pharmacist or clinician. Ask:

Keep a current list of medicines, vitamins, herbs, and supplements. Include the brand, ingredient amount, and how often you take each one. This is particularly important if you use an antidepressant or have a history of bipolar disorder.

Do not stop prescribed treatment in order to try SAM-e without discussing the change with the clinician responsible for that treatment. Sudden changes to some psychiatric medicines can cause withdrawal effects or a return of symptoms.

If a clinician is investigating anaemia, nerve symptoms, poor nutrition, or raised homocysteine, they may order tests such as vitamin B12, folate, or homocysteine. These tests assess related parts of metabolism; they do not directly measure whether you need SAM-e.

- Whether it could interact with your medicines or other supplements
- Whether your reason for taking it needs a separate assessment
- What benefit you could reasonably expect and how you would track it
- When to stop if it does not help or causes unwanted effects

Common questions

Is SAM-e the same as methionine?

No. Methionine is an amino acid obtained from food and recycled within the body. Cells use methionine and ATP to make SAM-e.

Can food provide SAM-e?

SAM-e is mainly made inside your cells. Protein-containing foods provide methionine, but there is no established diet designed to raise SAM-e levels in people without a diagnosed nutritional problem.

Does SAM-e increase serotonin?

SAM-e participates in chemical pathways involved in brain signalling, but it is not serotonin. Its effects are more complex than directly adding serotonin to the body. Because it may increase serotonin activity, combining it with serotonin-affecting medicines can be risky.

Can a blood test show low SAM-e?

SAM-e can be measured in research or specialised laboratory settings, but testing is not part of routine clinical care. There is no widely accepted blood level used to diagnose a general SAM-e deficiency or decide who should take a supplement.

Is "natural" SAM-e safer than medicine?

Not necessarily. A substance your body naturally makes can still cause side effects when taken in a concentrated manufactured form. Supplements can also interact with medicines and may differ in quality.

Does SAM-e treat depression or arthritis?

It has been studied for both conditions, but current evidence does not establish it as a reliable replacement for standard care.[3,4] If mood or joint symptoms are affecting daily life, an assessment can identify the condition and the options with the clearest evidence.

Sources

- National Center for Complementary and Integrative Health. S-Adenosyl-L-Methionine (SAMe): In Depth . National Institutes of Health.
- US Food and Drug Administration. Questions and Answers on Dietary Supplements .
- Galizia I, Oldani L, Macritchie K, et al. S-adenosyl methionine for depression in adults. Cochrane Database of Systematic Reviews . 2016;(10):CD011286.
- Rutjes AWS, Nüesch E, Reichenbach S, Jüni P. S-adenosylmethionine for osteoarthritis of the knee or hip. Cochrane Database of Systematic Reviews . 2009;(4):CD007321.
- Lu SC, Mato JM. S-adenosylmethionine in liver health, injury, and cancer. Physiological Reviews . 2012;92(4):1515-1542.
- Brosnan JT, Brosnan ME. The sulfur-containing amino acids: an overview. Journal of Nutrition . 2006;136(6 Suppl):1636S-1640S.