HEALTH BASICS

Subcutaneous

Subcutaneous means beneath the skin, usually referring to a tissue layer or a route for giving medicine.

WHAT IT IS
Tissue or a location directly beneath the skin

WHERE IT COMES UP
Injections, scans, examination notes, and descriptions of lumps

WHY IT MATTERS
It affects medicine delivery and interpretation of findings

WHAT IT IS NOT
Not a disease or diagnosis

WHO TO ASK
Pharmacist, nurse, or clinician who ordered the test

HOW IT IS CHECKED
Examination or imaging for a specific concern

Overview

Subcutaneous means under the skin. Clinicians may use the word to describe a layer of tissue, the location of a finding, or a way of giving medicine.

The subcutaneous layer lies below the dermis, which is the deeper part of the skin. It is also called subcutaneous tissue or the hypodermis. It contains fat, loose supporting tissue, blood vessels, and nerves. Its thickness differs across the body and from one person to another.

On a scan or examination note, "subcutaneous" usually tells you where something is. For example, a report may describe a subcutaneous lump, swelling, fluid collection, or area of inflammation. The word alone does not say what caused the finding or whether it is harmful.

A subcutaneous injection places medicine into this tissue rather than into a vein or muscle. Insulin, some blood-thinning medicines, and some other injected medicines are given this way.

How it works

Subcutaneous tissue connects the skin to structures underneath it. Its fat helps cushion the

body, reduce heat loss, and store energy. The tissue also allows the skin to move over muscles and other deeper structures.

Blood vessels pass through this layer, but it has less blood flow than muscle. Medicine injected there is absorbed through small blood vessels and lymph vessels. Absorption is generally slower than direct injection into a vein, but the exact pattern is specific to the medicine and its formulation.

Several factors can affect absorption. These include the injection site, local blood flow, temperature, exercise, and changes in the tissue from repeated injections. This is one reason the instructions for one medicine should not be used for another.

A needle intended for a subcutaneous injection should reach the tissue beneath the skin without entering muscle. Needle length, device design, and technique differ among medicines and people. Prefilled pens, syringes, pumps, and other devices may each have their own instructions.

"Subcutaneous" is different from two other common injection terms:

These routes are not interchangeable. Changing the route can alter how quickly a medicine works and may make treatment unsafe or ineffective.

- Intradermal means within the skin, as with certain skin tests
- Intramuscular means within a muscle, as with many vaccines
- Intravenous means directly into a vein

Why it matters for your health

The meaning of "subcutaneous" depends on its context.

For medicine, the word identifies the route of administration. Using the correct route helps the medicine reach the body in the way it was designed to. A label reading "for subcutaneous use" does not mean the medicine can also be injected into a muscle or vein.

For a lump or scan finding, the word identifies its depth. Knowing that a finding is in the subcutaneous tissue can help a clinician narrow down possible causes and decide whether examination, monitoring, ultrasound, or other imaging is useful.

Many different things can occur in this layer. A lipoma is a usually noncancerous growth made of fat cells. A cyst is a sac that may contain fluid or thicker material. A hematoma is a collection of blood, often after an injury or procedure. Infection may produce inflammation or an abscess, which is a collection of pus.

Subcutaneous tissue can also hold excess fluid, called edema. Air trapped in the tissue is called

subcutaneous emphysema. Each of these is a separate finding or condition. They are not meanings of the word "subcutaneous" itself.

What can go wrong

Repeated injections can irritate the skin and underlying tissue. Bruising, mild soreness, or a small amount of bleeding may occur around an injection site.

Repeated insulin injections into the same area can cause lipohypertrophy , meaning thickened or lumpy fatty tissue. Injecting into this altered tissue can make insulin absorption less predictable. Less commonly, fat tissue may shrink in an area, which is called lipoatrophy.

An injection site can become infected. Cellulitis is an infection of the deeper skin and subcutaneous tissue. An abscess is a pocket of infection that may need drainage. Spreading redness, increasing warmth, pus, fever, or worsening pain should be assessed promptly rather than left until a routine appointment.

A medicine can also cause an allergic reaction, regardless of where it is injected. Call emergency services for trouble breathing, wheezing, faintness, or swelling of the tongue or throat after an injection.

Subcutaneous lumps have many possible causes. A clinician may consider a lipoma, cyst, inflamed lymph node, hematoma, abscess, medication-related tissue change, or soft-tissue tumour. The location alone cannot distinguish among them. A lump that is growing, painful, firm, fixed in place, or persistent deserves clinical assessment, but these features do not establish a diagnosis by themselves.

Subcutaneous emphysema can occur after chest or neck injury, certain medical procedures, lung problems, or, rarely, serious infection. The area may feel crackly when touched because of air beneath the skin. Sudden neck or facial swelling, breathing difficulty, chest pain, voice changes, or rapid worsening requires emergency care. New crackling swelling after an injury or procedure should be reported promptly even if breathing is normal.

What you can do

You do not need a routine test simply because the word "subcutaneous" appears in a record. Ask what structure or finding the word describes and whether any follow-up is recommended.

If you give yourself a subcutaneous medicine, follow the instructions supplied for that specific product. Ask a pharmacist or nurse to demonstrate the device if you are uncertain. Do not change the injection route, needle, dose, or device based on general online instructions.

Useful steps include:

If you find a lump, note when you first noticed it, where it is, and whether it changes. A photograph or simple size record may help with routine follow-up. Do not squeeze, puncture, or try to drain an unexplained lump yourself.

Arrange routine assessment for a persistent or changing lump, repeated injection-site problems, or questions about a scan report. Seek prompt care for spreading redness, pus, fever, or worsening pain. Use emergency services for breathing difficulty, severe allergic-reaction signs, chest pain, or rapidly increasing swelling of the face or neck.

- Confirming the medicine is intended for subcutaneous use
- Following its specific instructions for approved injection areas
- Rotating sites as instructed rather than repeatedly using one spot
- Checking the skin for lumps, broken skin, or increasing redness
- Recording unexplained bruising or repeated injection problems
- Placing used needles directly into an approved sharps container
- Never sharing or reusing needles, syringes, or injection pens

Common questions

Is subcutaneous tissue the same as body fat?

Not exactly. Fat makes up much of this tissue, but the layer also contains connective tissue, blood vessels, nerves, and lymph vessels.

Does "subcutaneous" on a scan mean cancer?

No. It only identifies the location beneath the skin. The rest of the report, the examination, and sometimes further imaging help explain what the finding may be.

Is a subcutaneous injection always less painful than a muscle injection?

Not necessarily. Comfort can be affected by the medicine, device, injection site, technique, and the condition of the tissue.

Can I use any fatty area for an injection?

No. Approved sites differ by medicine and device. Use the product instructions and the technique shown by your pharmacist, nurse, or prescribing service.

Why should injection sites be rotated?

Repeatedly using one small area can damage or thicken the tissue. For medicines such as

insulin, this may also make absorption less predictable.

How is a subcutaneous lump checked?

A clinician usually begins by examining it and asking how long it has been present. Ultrasound can show whether a lump is solid or fluid-filled and how it relates to nearby tissue. Other imaging or a biopsy, which removes cells or tissue for testing, may be considered for selected findings.

Sources

- National Cancer Institute. NCI Dictionary of Cancer Terms: Subcutaneous . National Institutes of Health.
- OpenStax. Anatomy and Physiology 2e, Section 5.1: Layers of the Skin . Rice University.
- Centers for Disease Control and Prevention. Preventing Unsafe Injection Practices . Injection Safety.
- US Food and Drug Administration. Best Way to Get Rid of Used Needles and Other Sharps .
- Frid AH, Kreugel G, Grassi G, et al. "New Insulin Delivery Recommendations." Mayo Clinic Proceedings . 2016;91(9):1231-1255.
- American College of Radiology. ACR Appropriateness Criteria: Soft Tissue Masses . 2022.
- Kukuruza K, Aboeed A. "Subcutaneous Emphysema." StatPearls . National Center for Biotechnology Information Bookshelf.