HEALTH BASICS

Swallowing gum

Swallowing gum means chewing gum enters the digestive tract instead of being spat out.

WHAT IT IS
Chewing gum that has been swallowed

WHERE IT COMES UP
After accidental swallowing, often in children

WHY IT MATTERS
Usually harmless; rare choking or bowel blockage

WHAT IT IS NOT
Not a disease or diagnosis

WHO TO ASK
Poison centre or urgent care for concerning symptoms

HOW IT IS CHECKED
None routinely; imaging or endoscopy if blockage is suspected

Overview

Swallowing gum is not a disease or diagnosis. It is usually a one-time accident, although some people swallow gum out of habit.

A common myth says swallowed gum stays in the stomach for seven years. It does not. The body cannot fully break down the gum base, but the digestive tract usually moves it along with other material. It then leaves the body in stool.

A single piece of ordinary chewing gum generally does not require a test, treatment, or follow-up if you feel well. Swallowing a large amount, repeatedly swallowing gum, or swallowing medicated gum may need separate advice.

The immediate concern is different if the gum enters the airway instead of the digestive tract. Gum in the airway can cause choking, particularly in a young child.

How it works

Chewing gum usually contains a gum base, flavourings, sweeteners, colours, and ingredients that keep it soft. Your saliva dissolves some of these substances while you chew.

When gum is swallowed, it passes from the mouth through the oesophagus, the tube that carries food to the stomach. The stomach and small intestine absorb ingredients they can break down.

The gum base is different. Digestive acids and enzymes cannot fully digest it. However, "not digestible" does not mean that it remains in the stomach. Normal muscular contractions move the gum base through the intestines with food, fluid, and other waste.

You may not notice it in your stool. There is usually no reason to look for it, and its appearance cannot reliably show whether everything is normal.

Sugar-free gum often contains sugar alcohols, such as sorbitol or xylitol. These can cause gas, cramping, or loose stools when consumed in larger amounts. This effect comes from the sweetener rather than from gum becoming stuck.

Why it matters for your health

For someone who has swallowed one piece of ordinary gum and has no concerning symptoms, the health effect is usually minimal. No special food, drink, or medicine is needed to "push" the gum through.

The amount and circumstances matter. Repeatedly swallowing many pieces can allow gum to collect with other material. This is uncommon, but it has been reported, especially in children with constipation.

The type of gum also matters. Nicotine gum and other medicated gums contain active drugs. Their packaging includes instructions about how they should be used. Swallowing one of these products should be treated as a possible medication exposure, not simply as swallowed ordinary gum.

Swallowing gum during fasting for surgery or a medical procedure is another special situation. Tell the procedure or anaesthesia team what happened and when. Do not decide on your own whether the procedure should continue, because fasting instructions differ by procedure and facility.

What can go wrong

Choking happens when gum blocks the airway. This is an emergency if the person cannot breathe, speak, or cough effectively, becomes blue or grey, or collapses. Call emergency services and follow their choking instructions. Use first aid only if you know the correct method for the person's age and situation.

Oesophageal obstruction means something is stuck in the tube between the mouth and stomach. Seek urgent medical help if the person cannot swallow saliva, is drooling continuously, has persistent chest or throat pain, or repeatedly brings food and drink back up.

A bezoar is a firm mass of material that has collected in the digestive tract. Rare reports describe chewing-gum bezoars after children repeatedly swallowed gum, sometimes together with constipation or other swallowed material. A bezoar can contribute to bowel obstruction , meaning the intestine is partly or completely blocked.

Get urgent medical advice for severe or worsening abdominal pain, repeated vomiting, a swollen abdomen, blood in vomit or stool, or inability to pass stool or gas. These signs have many possible causes and do not prove that gum is responsible.

Contact a poison centre or urgent medical service if a large or uncertain amount was swallowed, if the gum contained nicotine or another medicine, or if other substances were swallowed with it. Have the package available if possible.

What you can do

If you swallowed one piece of ordinary gum and feel well, there is usually nothing specific you need to do. You can continue your normal meals and fluids.

Do not make yourself vomit. Do not take a laxative just because you swallowed gum. These actions are unlikely to help and can cause other problems.

If advice is needed, keep a brief record of:

For a child, check whether anything else is missing, such as the wrapper, packaging, coins, magnets, or batteries. Button batteries and magnets require urgent specialist advice because they can seriously injure the digestive tract.

If a child repeatedly eats or swallows non-food items, mention this at a routine healthcare visit. The clinician may ask about eating habits, development, constipation, and pica , which means repeatedly eating items that are not food.

Children should only have gum when they can reliably chew it without choking and understand that it should be spat out. Gum should not be given while a child is running, playing actively, or lying down.

- The product name and whether it was medicated
- The approximate number of pieces
- The time it happened

- Any symptoms that followed
- Relevant issues such as constipation or previous intestinal surgery

Common questions

Does gum stay in your stomach for seven years?

No. The gum base is not fully digested, but it usually travels through the digestive tract and leaves in stool. It does not normally attach to the stomach for years.

Can stomach acid dissolve gum?

Stomach acid can help release some ingredients, but it does not fully break down the gum base. The intestines can still move that base along.

Should I eat bread or another food to move it through?

No special food has been shown to be necessary. If you feel well, continue your usual diet rather than trying to force the gum through.

Do I need to check my stool?

Usually not. Gum may be difficult to recognise after passing through the digestive tract, and finding it does not provide useful health information.

Is swallowing sugar-free gum different?

The gum base follows the same general path. However, consuming a large amount of certain sugar alcohol sweeteners can cause gas, cramps, or diarrhoea.

What if I swallowed nicotine gum?

Nicotine gum is a medicine. Contact a poison centre or healthcare service for advice based on the product, amount, age, and symptoms. Keep the packaging nearby. Seek emergency help for collapse, trouble breathing, a seizure, or severe confusion.

When is routine follow-up needed?

A single symptom-free episode usually needs no follow-up. Arrange a routine appointment if swallowing gum happens often, is linked with ongoing constipation, or is part of repeatedly swallowing non-food items.

Sources

- Mayo Clinic Staff. "Swallowing gum: Is it harmful?" Mayo Clinic Expert Answers .

- Milov DE, Andres JM, Erhart NA, Bailey DJ. "Chewing Gum Bezoars of the Gastrointestinal Tract." Pediatrics . 1998;102(2):e22. doi:10.1542/peds.102.2.e22.
- Kramer RE, Lerner DG, Lin T, et al. "Management of Ingested Foreign Bodies in Children: A Clinical Report of the North American Society for Pediatric Gastroenterology, Hepatology and Nutrition Endoscopy Committee." Journal of Pediatric Gastroenterology and Nutrition . 2015;60(4):562-574. doi:10.1097/MPG.0000000000000729.
- MSD Manual Consumer Version. "Bezoars." Reviewed by Jonathan Gotfried, MD.
- National Library of Medicine. DailyMed: Nicorette nicotine polacrilex gum-drug label .