CONDITION

Lymphangitis

Lymphangitis is inflammation of lymph vessels, usually caused by a bacterial skin infection.

WHAT IT IS
Inflammation of lymph vessels, usually from bacterial skin infection

BODY SYSTEM
Lymphatic and immune systems

MAIN SYMPTOMS
Red streaks, skin pain, tender lymph nodes, fever

WHAT CAUSES IT
Usually bacterial infection entering through damaged skin

HOW IT IS DIAGNOSED
Clinical examination; cultures for selected wounds or severe illness

MAIN TREATMENT
Antibiotics, wound care, and treatment of any abscess

CAN IT BE CURED?
Yes - antibiotics usually clear infection

IS IT AN EMERGENCY?
Sometimes - rapid spread or sepsis symptoms

LONG-TERM OUTLOOK
Most people recover fully with prompt treatment

Overview

Lymphangitis is inflammation of the lymph vessels. These small channels carry lymph, a clear fluid containing infection-fighting white blood cells, through the body.

Most cases happen when bacteria enter damaged skin and spread into nearby lymph vessels. A painful red or darker streak may extend from a wound or infected area toward a group of lymph nodes, such as those in the armpit or groin.

Lymphangitis often develops alongside cellulitis, which is an infection of the deeper layers of skin. It can also occur with an abscess, an enclosed pocket of pus.

The condition matters because infection can spread quickly through the tissues and, less commonly, into the bloodstream. Prompt assessment and antibiotic treatment usually lead to a full recovery. A red streak does not automatically mean there is a bloodstream infection, but

it should not be ignored.

Less common forms develop more slowly and may cause a line of firm bumps rather than a smooth red streak. These can be linked to certain fungi or unusual bacteria and need different treatment.

Symptoms

Symptoms of acute bacterial lymphangitis may develop over several hours or days. They usually begin near a cut, bite, blister, ulcer, or other area of skin infection.

Possible symptoms include:

The streak usually follows the path of a lymph vessel toward nearby lymph nodes. On brown or black skin, redness may be less obvious. The affected area may instead look darker, purple, or grey, or may be easier to identify by warmth, swelling, and tenderness.

A slower form called nodular lymphangitis causes a chain of firm bumps under or within the skin. Some bumps may break open. This pattern can develop over days or weeks after contact with soil, plants, animals, or contaminated water.

Not every red line on the skin is lymphangitis. Irritation, an inflamed vein, an allergic reaction, or another skin condition may look similar. A healthcare professional can assess the pattern and other symptoms.

- A red, pink, purple, or darker streak moving away from the affected skin
- Warmth, swelling, tenderness, or throbbing pain
- Painful or enlarged lymph nodes, often in the armpit, groin, or neck
- Fever, chills, headache, tiredness, or muscle aches
- Pus, fluid, or a crusted area at the original wound

Causes and risk factors

Acute lymphangitis is usually caused by bacteria entering through a break in the skin. Group A Streptococcus is a common cause. Staphylococcus aureus and other bacteria can also be responsible.

The opening may be easy to see, such as a cut or puncture. Sometimes it is very small, including a crack between the toes caused by athlete's foot.

Possible entry points include:

Risk may be higher when lymph drainage or circulation is already reduced. This can happen with lymphoedema, which is long-term swelling caused by a build-up of lymph fluid. Previous lymph node surgery, vein problems, diabetes, and conditions or medicines that weaken immune defences can also increase risk.

Uncommon causes include fungi such as Sporothrix , which lives in soil and plant material, and certain bacteria associated with animals or water. These infections often produce slowly developing nodules rather than a rapidly spreading streak.

Lymphangitis itself is not inherited. It is also not usually passed directly from one person to another, although bacteria from an uncovered, draining wound may spread through close contact.

- Cuts, grazes, burns, splinters, or puncture wounds
- Insect, animal, or human bites
- Blisters, eczema, fungal infections, or cracked skin
- Leg ulcers or surgical wounds
- Injected substances or medical devices that pass through the skin

Diagnosis

Lymphangitis is usually diagnosed from your symptoms and a physical examination. There is no single blood test or scan that confirms every case.

A healthcare professional may examine the original wound, the direction of any streak, nearby lymph nodes, and the extent of warmth and swelling. They may also check your temperature, heart rate, breathing, and blood pressure for signs that the infection is affecting your whole body.

If there is pus or an open wound, a sample may be sent to a laboratory. This can identify the organism and help guide treatment, especially if the infection is severe, unusual, or not improving.

Blood tests may show inflammation but do not prove that lymphangitis is present. Blood cultures, which look for bacteria in the bloodstream, are generally reserved for people with significant whole-body symptoms or particular medical risks.

Ultrasound or other imaging is not routinely needed. It may be used if there could be an abscess, a blood clot, a foreign object, or a deeper infection.

Tell the healthcare professional about recent bites, gardening injuries, animal contact, water

exposure, travel, injected substances, and medicines that affect your immune system. These details can point to less common causes.

Treatment and management

Bacterial lymphangitis is treated with antibiotics like amoxicillin or doxycycline . The choice and route depend on how severe the infection is, its likely source, local bacterial patterns, allergies, and other health conditions.

People who are otherwise well may be treated with antibiotic tablets. Hospital care and antibiotics given into a vein may be needed if the infection is spreading quickly, whole-body symptoms are significant, tablets cannot be taken, or immune defences are reduced.

If there is an abscess, antibiotics may not be enough by themselves. A healthcare professional may need to drain it. Any splinter, dead tissue, infected device, or other continuing source may also need attention.

Supportive care may include:

Do not squeeze, cut, or attempt to drain the area yourself. Take prescribed antibiotics exactly as directed and complete the course unless the prescriber tells you to stop or change it.

Fungal and unusual bacterial causes require different medicines, often for longer periods. Standard antibiotics used for routine skin infections may not work for them. Laboratory testing or review by a specialist may therefore be needed when nodules develop slowly or initial treatment fails.

- Resting and raising the affected limb to reduce swelling
- Using a warm, clean compress for comfort
- Drinking enough fluid, especially when fever is present
- Using suitable non-prescription pain or fever medicine as directed
- Keeping the original wound clean and covered

Complications

Most people recover without lasting problems. Complications are more likely when treatment is delayed, the infection is severe, or immune defences and lymph drainage are reduced.

Possible complications include an abscess, worsening cellulitis, or bacteria entering the bloodstream. A widespread response to infection is called sepsis and needs emergency

treatment.

Rarely, a deeper, rapidly progressing tissue infection can develop. Severe pain that seems greater than the visible skin changes, blistering, skin turning dusky or black, or very rapid spread requires emergency assessment.

Repeated or severe infections can damage lymph vessels. This may contribute to ongoing swelling and make future skin infections more likely.

Prevention

Not every case can be prevented. Small skin breaks may go unnoticed, and lymphangitis can occasionally develop despite good wound care.

You can reduce the chance of bacteria entering the skin by washing cuts and grazes with clean water, covering them with a clean dressing, and watching for increasing pain, warmth, swelling, or discharge. Avoid picking at wounds or sharing equipment that pierces the skin.

Treating cracked skin, eczema, athlete's foot, and ulcers can help protect the skin barrier. Moisturising dry skin may reduce splitting, but moisturiser should not be placed inside an open wound.

If you have diabetes, poor circulation, or lymphoedema, regular skin and foot checks can help you notice injuries early. Follow any individual advice you have been given about swelling management or protecting a limb after lymph node treatment.

Animal or human bites, deep punctures, and dirty wounds may need medical assessment. A healthcare professional can also check whether tetanus protection is up to date.

Living with it and what to expect

Acute bacterial lymphangitis is usually a short-term condition. With effective treatment, fever and spreading inflammation often begin to settle within a day or two, although skin tenderness and colour changes may take longer to disappear. Recovery is not identical for everyone.

The affected streak should stop extending. Pain, warmth, and swelling should gradually ease. A clinician may mark the edge of the affected area to make changes easier to see.

If symptoms continue to spread, remain unchanged after treatment has had time to work, or return soon afterward, further assessment is needed. The bacteria may not be covered by the first antibiotic, or an abscess or foreign object may be present. A slower fungal or unusual bacterial infection may also need to be considered.

Most people do not have permanent effects after one promptly treated episode. Recurrence is more likely when there is ongoing lymphoedema, repeated skin damage, chronic ulcers, or an untreated fungal infection between the toes. Managing these underlying problems can lower future risk.

Nodular lymphangitis caused by fungi or unusual bacteria may last for weeks or months before the cause is identified. Recovery usually takes longer because these infections need specialised treatment.

When to get medical help

Seek same-day medical advice if you develop a tender red or dark streak extending from a wound, especially if it is spreading or you also have fever, chills, swollen lymph nodes, or increasing pain. Prompt assessment is also important if you have diabetes, lymphoedema, poor circulation, or weakened immune defences.

Seek emergency care if you have possible signs of sepsis or a rapidly progressing deep infection, including:

Arrange routine follow-up if symptoms are not beginning to improve after the timeframe given by the treating healthcare professional, if they worsen at any point, or if lymphangitis keeps returning. New diarrhoea, rash, vomiting, or other possible antibiotic side effects should also be reported.

- Confusion, fainting, severe drowsiness, or difficulty waking
- Trouble breathing or very rapid breathing
- Cold, clammy, mottled, or unusually pale skin
- Severe weakness or a feeling that something is seriously wrong
- Rapidly spreading swelling, severe pain, blisters, or dusky skin
- Signs of a severe allergic reaction after medicine, such as facial swelling or breathing difficulty

Common questions

Is lymphangitis the same as blood poisoning?

No. Lymphangitis affects lymph vessels, not blood vessels. However, the same infection can sometimes enter the bloodstream and lead to sepsis. A red streak alone does not prove that

this has happened.

Can lymphangitis go away without antibiotics?

Bacterial lymphangitis generally needs prompt antibiotic treatment. Waiting for it to clear by itself risks further spread. Fungal or unusual bacterial forms also need treatment chosen for the specific cause.

Is lymphangitis contagious?

The inflammation in the lymph vessels is not contagious. Bacteria from an open or draining wound may spread through direct contact, so keep the area covered, wash your hands after touching the dressing, and do not share towels.

How quickly does lymphangitis spread?

Acute bacterial lymphangitis can extend over hours or days. The speed differs according to the organism, the wound, immune defences, circulation, and whether treatment has started. A visibly advancing streak needs same-day assessment.

What is the difference between lymphangitis and cellulitis?

Cellulitis is infection and inflammation within the deeper skin and tissue beneath it. Lymphangitis affects the lymph vessels and often appears as a streak leading away from cellulitis or a wound. Both can occur at the same time.

Does every red line from a wound mean lymphangitis?

No. Skin irritation, an inflamed vein, scratching, and allergic reactions can also cause lines or redness. Warmth, tenderness, spreading colour change, swollen lymph nodes, and fever make infection more concerning, but an examination is needed to identify the cause.

Sources

- Centers for Disease Control and Prevention. "Clinical Guidance for Group A Streptococcal Cellulitis."
- MedlinePlus Medical Encyclopedia, US National Library of Medicine. "Lymphangitis."
- Merck Manual Professional Edition. "Lymphangitis."
- Stevens DL, Bisno AL, Chambers HF, et al. "Practice Guidelines for the Diagnosis and Management of Skin and Soft Tissue Infections: 2014 Update by the Infectious Diseases Society of America." Clinical Infectious Diseases . 2014;59(2):e10-e52.
- Tirado-Sánchez A, Bonifaz A. "Nodular Lymphangitis (Sporotrichoid Lymphocutaneous Infections): Clues to Differential Diagnosis." Journal of Fungi . 2018;4(2):56.