CONDITION

Macroglossia

Macroglossia is an unusually enlarged tongue that may affect breathing, eating, speech, or dental development.

WHAT IT IS
Persistent tongue enlargement beyond the mouth's usual proportions

BODY SYSTEM
Mouth, upper airway, and digestive system

MAIN SYMPTOMS
Tongue protrusion, drooling, feeding difficulty, speech changes

WHAT CAUSES IT
Inherited conditions, tissue overgrowth, swelling, or acquired disease

HOW IT IS DIAGNOSED
Clinical examination, plus tests to identify the cause

MAIN TREATMENT
Cause-specific care, therapy, dental care, or reduction surgery

CAN IT BE CURED?
Sometimes - treatable causes or surgery can reduce enlargement

IS IT AN EMERGENCY?
Sometimes - breathing difficulty requires emergency care

LONG-TERM OUTLOOK
Often manageable; outlook reflects airway problems and underlying cause

Overview

Macroglossia means that the tongue is larger than expected for the size of the mouth. The enlargement may involve the whole tongue or only one area. It can be present at birth, develop during childhood, or appear later in life.

Some people have true macroglossia , in which the tongue tissue itself is enlarged. Others have relative macroglossia , in which a normal-sized tongue seems too large because the jaw or mouth is small.

Macroglossia matters when the tongue interferes with breathing, eating, swallowing, speech, sleep, or dental development. A tongue that repeatedly rests outside the mouth can also become dry, cracked, injured, or infected.

The condition ranges from mild to severe. A mildly enlarged tongue may need only monitoring and supportive care. More significant enlargement may require treatment of an underlying condition, feeding or speech support, dental care, or surgery.

Sudden tongue swelling is different from long-standing macroglossia. Rapid swelling can narrow the airway and needs urgent assessment, especially if breathing or swallowing is difficult.

Symptoms

The appearance and effects of macroglossia differ from person to person. The tongue may fill most of the mouth, press against the teeth, or protrude between the lips.

Possible symptoms and signs include:

Babies may have difficulty attaching to the breast or bottle, coordinating sucking and swallowing, or gaining weight. However, feeding problems have many possible causes and do not by themselves confirm macroglossia.

In older children, a persistently enlarged tongue can influence speech and the way the teeth and jaws develop. Adults may first notice tongue enlargement through snoring, dental impressions, discomfort, or increasing difficulty speaking or swallowing.

- A tongue that often rests outside the mouth
- Drooling or difficulty keeping saliva in the mouth
- Trouble sucking, chewing, moving food, or swallowing
- Speech that is difficult to understand
- Noisy breathing, snoring, or pauses in breathing during sleep
- Tooth marks or a scalloped pattern along the tongue's edges
- Mouth breathing or difficulty closing the lips
- Repeated tongue biting, sores, bleeding, or dryness
- Teeth that do not meet normally, called malocclusion
- Changes in jaw growth or spacing between the teeth

Causes and risk factors

Macroglossia is a physical finding rather than a single disease. Identifying its cause is important

because treatment and long-term outlook are linked to the underlying condition.

Some people are born with tongue enlargement. Congenital causes include tissue overgrowth and vascular or lymphatic malformations. These are abnormal collections of blood vessels or lymph channels. They may affect one part of the tongue and can sometimes swell after infection, injury, or bleeding.

Macroglossia can occur as part of an inherited or genetic condition. Examples include Beckwith-Wiedemann syndrome, Down syndrome, and some rare conditions that affect how the body stores or breaks down certain substances. In Down syndrome, relative macroglossia may be related partly to the size and shape of the mouth rather than major enlargement of the tongue itself.

Causes that can develop later include:

A suddenly swollen tongue may result from an allergic reaction or angioedema, which is deeper swelling beneath the skin and lining of the mouth. This is not the same as chronic macroglossia, but it can look similar and may become an emergency.

Sometimes no single cause is found.

- Amyloidosis , in which abnormal proteins build up in tissues
- Acromegaly , caused by excess growth hormone in adulthood
- Severe or untreated hypothyroidism , meaning an underactive thyroid
- Benign or cancerous growths involving the tongue
- Inflammation, infection, injury, bleeding, or fluid buildup
- Enlargement of tongue muscle or other local tissues

Diagnosis

Macroglossia is diagnosed mainly by examining the tongue, mouth, teeth, jaw, and airway. There is no universal measurement that confirms the condition in every person. The key question is whether the tongue is unusually large for the mouth and whether it affects function.

The assessment may include questions about when the enlargement began, whether it is changing, and whether there are problems with feeding, swallowing, speech, sleep, or breathing. In children, growth and dental development may also be reviewed.

Tests are selected according to the suspected cause. They may include:

Photographs, dental models, or imaging may help monitor changes over time. An examination

may involve specialists in ear, nose and throat care, oral and facial surgery, genetics, dentistry, speech and language therapy, or paediatrics.

- Blood tests for thyroid function, hormone disorders, or other illnesses
- Ultrasound or magnetic resonance imaging to examine the tongue's tissues
- A sleep study if snoring or pauses in breathing suggest sleep apnoea
- Swallowing or feeding assessment
- Hearing and speech assessment
- Genetic testing when an inherited condition is suspected
- Biopsy, which removes a small tissue sample, if a mass or amyloidosis is possible

Treatment and management

Treatment is based on the cause, the degree of enlargement, and the problems it creates. An enlarged tongue that is stable and does not affect breathing, feeding, speech, sleep, or dental development may only need routine monitoring.

When an underlying condition can be treated, doing so may reduce enlargement or prevent further change. For example, care may focus on a thyroid or hormone disorder. Vascular and lymphatic malformations may need specialist procedures or other targeted treatment.

Supportive care can help with day-to-day function. It may include:

Some children improve in function as the mouth and jaw grow, although this cannot be assumed for every child. Speech difficulties may relate to several factors, so reducing tongue size does not always correct speech without therapy.

A reduction glossectomy is surgery to remove part of the tongue and reshape it. It may be considered when macroglossia causes ongoing airway obstruction, serious feeding problems, repeated injury, major speech difficulties, or harmful changes to the teeth and jaws.

Surgery is planned carefully because the tongue is important for taste, speech, swallowing, and airway protection. Possible risks include bleeding, swelling, infection, scarring, altered sensation, changes in movement, and the need for further treatment. The timing and likely benefit depend on the cause and pattern of enlargement.

- Feeding advice and swallowing therapy
- Speech and language therapy
- Dental or orthodontic monitoring

- Care for dry, cracked, or repeatedly injured tongue tissue
- Sleep assessment and treatment for sleep-related breathing problems

Complications

Macroglossia can narrow the upper airway, especially during sleep. This may contribute to obstructive sleep apnoea, in which breathing repeatedly becomes reduced or stops for short periods.

Feeding and swallowing problems can make meals tiring and may affect nutrition or growth. If food or liquid enters the airway instead of the food pipe, it is called aspiration. Repeated aspiration can irritate the lungs and contribute to chest infections.

Pressure from the tongue may gradually move the teeth or affect jaw development. Possible changes include gaps between teeth, an open bite in which the front teeth do not meet, or a lower jaw that appears more prominent.

A tongue that remains outside the mouth may become dry, cracked, swollen, or injured. Repeated biting can cause ulcers and bleeding. Drooling and visible tongue protrusion may also affect comfort, communication, and social confidence.

The underlying cause may have complications separate from the tongue enlargement. Follow-up therefore often includes monitoring the broader condition, not only the tongue.

Prevention

Many cases of macroglossia cannot be prevented. This is particularly true when it is caused by an inherited condition, congenital tissue overgrowth, or a vascular or lymphatic malformation.

Prompt treatment of certain acquired conditions may limit progression, but it does not guarantee that tongue enlargement will reverse. Following the monitoring plan for a known thyroid, hormone, metabolic, or protein-deposition disorder may help identify changes earlier.

Protecting a protruding tongue from repeated biting, excessive dryness, and sharp dental edges may reduce injury. This manages complications rather than preventing macroglossia itself.

Living with it and what to expect

The course of macroglossia depends mainly on its cause. Some congenital forms remain fairly

stable. Others become more noticeable as a child grows or swell at certain times. Enlargement linked to an acquired illness may improve when that illness is treated, remain unchanged, or progress.

Regular follow-up may track breathing, sleep, nutrition, speech, tongue injury, and dental development. Children may need reassessment as their face and jaws grow. Adults with a new or enlarging tongue may need continued evaluation for an underlying illness.

Many people with mild macroglossia live without major restrictions. Those with feeding, speech, sleep, or dental problems may need care from several specialties over months or years. Therapy and dental treatment can improve function even when the tongue remains enlarged.

Surgery can reduce tongue size and relieve significant functional problems, but it does not always remove every symptom. Further dental work, speech therapy, or treatment for the underlying condition may still be needed. Some causes can also lead to renewed enlargement after treatment.

When to get medical help

Call emergency services if tongue swelling is sudden or if there is:

Do not wait for a routine appointment when breathing is affected. Rapid swelling can progress quickly.

Seek prompt medical assessment if an enlarged tongue is accompanied by choking, repeated coughing during meals, poor feeding, dehydration, unexplained weight loss, repeated chest infections, or pauses in breathing during sleep.

Arrange routine follow-up for a tongue that appears persistently large, is gradually enlarging, rests outside the mouth, or causes speech, dental, swallowing, or sleep problems. New tongue enlargement in an adult should also be assessed, even when it is painless.

- Difficulty breathing, gasping, or worsening noisy breathing
- An inability to swallow saliva
- Swelling of the lips, face, or throat
- Faintness, widespread hives, or other signs of a severe allergic reaction
- Blue, grey, or unusually pale lips or skin

Common questions

Can macroglossia go away on its own?

Sometimes the appearance becomes less noticeable as a child's mouth and jaw grow. Enlargement caused by a treatable illness may improve with treatment. Congenital tissue overgrowth and malformations often persist, although their effects can be managed.

Is macroglossia cancer?

Macroglossia itself is not cancer. Rarely, a growth in the tongue can make it larger. A new lump, one-sided enlargement, unexplained bleeding, persistent pain, or rapid change needs medical assessment.

Can a baby be born with macroglossia?

Yes. It may occur alone or as part of a congenital or genetic condition. A baby with a large tongue may need assessment of feeding, breathing, growth, and other physical features.

Does macroglossia affect speech and eating?

It can. The tongue may have less room to move food or form speech sounds. The amount of difficulty does not always match the tongue's visible size, so feeding, swallowing, and speech assessments can be useful.

Does a scalloped tongue mean I have macroglossia?

No. Tooth marks can appear when the tongue presses against the teeth, but they can also occur without true enlargement. The whole mouth, jaw, symptoms, and medical history need to be considered.

Is tongue-reduction surgery always needed?

No. Surgery is generally reserved for significant problems such as airway obstruction, serious feeding difficulty, repeated injury, or harmful dental and jaw changes. Mild, stable macroglossia may be monitored without surgery.

Sources

- Topouzelis N, Iliopoulos C, Kolokitha OE. "Macroglossia." International Dental Journal . 2011;61(2):63-69.
- National Center for Biotechnology Information. "Macroglossia." StatPearls . StatPearls Publishing.
- Shuman C, Kalish JM, Weksberg R. "Beckwith-Wiedemann Syndrome." GeneReviews . University of Washington, Seattle.

- International Society for the Study of Vascular Anomalies. ISSVA Classification for Vascular Anomalies .
- National Institute of Diabetes and Digestive and Kidney Diseases. "Acromegaly." National Institutes of Health.
- National Health Service. "Amyloidosis." NHS.