CONDITION

Necrotizing granuloma

A necrotizing granuloma is an area of organized inflammation containing damaged or dead tissue.

WHAT IT IS
Organized inflammation surrounding damaged or dead tissue

BODY SYSTEM
Lungs most often; sometimes other organs

MAIN SYMPTOMS
Often none, cough, fever, weight loss

WHAT CAUSES IT
Usually infection; sometimes immune inflammation or unknown

HOW IT IS DIAGNOSED
Tissue biopsy, microbial tests, and imaging

MAIN TREATMENT
Cause-specific antimicrobial or immune treatment

CAN IT BE CURED?
Sometimes - the underlying cause is treatable

IS IT AN EMERGENCY?
Sometimes - severe breathlessness or significant coughing of blood

LONG-TERM OUTLOOK
Often stable after the cause is treated

Overview

A necrotizing granuloma is a pattern seen when tissue is examined under a microscope. It is not one specific disease.

A granuloma is a small, organized collection of immune cells. Granulomas form when the immune system tries to contain something that is difficult to remove, such as certain germs or persistent inflammation. Necrotizing means that some tissue inside the granuloma has died or broken down.

Necrotizing granulomas are most often found in the lungs or nearby lymph nodes. They can also occur in the skin, digestive system, liver, spleen, bones, or other organs.

The finding matters because it can have several causes. Infections, particularly tuberculosis,

nontuberculous mycobacteria, and some fungal infections, are important possibilities. Less often, the cause is an inflammatory condition such as granulomatosis with polyangiitis, which affects small blood vessels. Sometimes testing does not identify a cause.

The word "necrotizing" does not mean that the tissue is cancerous. However, a lung granuloma can appear as a nodule or mass on a scan, so a biopsy may be needed to distinguish it from cancer and other conditions.

Symptoms

A necrotizing granuloma itself may not cause symptoms. Many lung granulomas are found by chance on an X-ray or CT scan performed for another reason.

When symptoms occur, they usually come from the underlying infection or inflammatory condition. A lung granuloma may be associated with:

Some people cough up blood. Small streaks can occur after forceful coughing, but this symptom still needs medical assessment.

Symptoms outside the lungs depend on the affected organ. A skin granuloma may cause a firm lump, sore, or ulcer. Enlarged lymph nodes may feel like painless or tender lumps. Involvement of the joints, kidneys, sinuses, or nerves can produce symptoms related to those areas.

Symptoms alone cannot show whether a granuloma is infectious, inflammatory, or inactive. Some infections cause few symptoms, while noninfectious inflammation can cause fever and weight loss that resemble an infection.

- A persistent cough
- Chest discomfort or shortness of breath
- Fever or night sweats
- Tiredness, reduced appetite, or unplanned weight loss

Causes and risk factors

Infections are a major cause of necrotizing granulomas. Important groups include:

The likely infection partly reflects where you have lived or travelled, environmental exposures, and how well your immune system is working. Some fungi are linked to particular geographic areas and are inhaled from soil or dust.

Noninfectious causes include granulomatosis with polyangiitis and some other forms of

blood-vessel inflammation. Rheumatoid nodules can also contain necrotizing granulomas. Rare inflammatory lung conditions may produce a similar pattern.

Factors that may influence the possible cause include:

These factors do not prove what caused a granuloma. People without an obvious risk factor can still develop one.

In some cases, no organism or inflammatory disease is found. This may happen because the original trigger has cleared, because only a small tissue sample was available, or because current tests cannot identify the cause.

- Mycobacteria , including the bacteria that cause tuberculosis and nontuberculous mycobacterial disease
- Fungi , such as organisms that cause histoplasmosis, coccidioidomycosis, blastomycosis, or cryptococcosis
- Less commonly, other bacteria or parasites
- Previous exposure to someone with tuberculosis
- Living or travelling in an area where certain fungal infections occur
- A weakened immune system from illness or immune-suppressing treatment
- Chronic lung disease
- Rheumatoid arthritis or symptoms of blood-vessel inflammation
- Work or activities involving substantial soil or dust exposure

Diagnosis

A CT scan can show the size, shape, location, and number of lung nodules. It may also show enlarged lymph nodes, cavities, or other lung changes. Imaging can suggest possible causes, but it usually cannot confirm a necrotizing granuloma.

Confirmation requires a biopsy , meaning that a sample of tissue is removed and examined under a microscope. Depending on the location, tissue may be collected during bronchoscopy, through a needle guided by a scan, during surgery, or from an accessible site such as the skin or a lymph node.

The pathologist looks at how the immune cells and necrotic tissue are arranged. Special stains may be used to search for mycobacteria and fungi. Tissue, mucus, or other samples may also be sent for:

A negative stain does not completely exclude infection. There may be too few organisms in the

sample to see. Cultures can add useful information, although some organisms grow slowly and some do not grow from the submitted sample.

The final interpretation combines the biopsy result with symptoms, imaging, exposure history, immune health, and microbiology results. Follow-up scans may be recommended when the cause remains uncertain or when a lung nodule has not been removed completely.

- Bacterial, mycobacterial, and fungal cultures
- Molecular tests that look for genetic material from an organism
- Tuberculosis blood or skin testing
- Blood or urine tests for specific fungal infections
- Tests for inflammation, kidney function, or immune-system antibodies

Treatment and management

Treatment is aimed at the underlying cause rather than at the granuloma alone.

Tuberculosis requires a combination of antimicrobial medicines and coordinated follow-up. Public health services may also help with testing close contacts. Nontuberculous mycobacterial disease is treated differently from tuberculosis. Treatment is not based on a positive culture alone; symptoms, scans, and repeated laboratory findings may all be considered.

Some fungal infections clear without treatment in otherwise healthy people. Others need antifungal medicine, especially when symptoms are significant, infection has spread, or the immune system is weakened. The organism and severity guide the treatment approach.

Inflammatory causes may be treated with corticosteroids or other medicines that reduce immune activity. Because these medicines can worsen an unrecognized infection, clinicians generally investigate infection before starting major immune-suppressing treatment.

Surgery may remove a solitary lung nodule when cancer cannot be excluded or when less invasive biopsy methods do not provide enough information. Surgery is not required for every granuloma.

No active treatment may be needed when a granuloma has been fully removed, testing does not suggest an active infection or inflammatory disease, and follow-up findings remain stable. Monitoring may include clinic visits, symptom review, and repeat imaging.

Do not stop immune-suppressing medicine or begin leftover antibiotics or antifungals like fluconazole or terbinafine on your own. The possible causes require different treatments, and

the wrong treatment can delay an accurate diagnosis.

Complications

Many isolated granulomas cause no lasting problem. Possible complications mainly reflect the underlying cause and the organ involved.

An active lung infection can cause cavities, scarring, or reduced lung function. It may spread beyond the lungs in people whose immune systems are weakened. Untreated tuberculosis can also spread to other people.

Blood-vessel inflammation can damage the lungs, kidneys, nerves, sinuses, or other organs. Early recognition is important because kidney inflammation may not cause obvious symptoms at first.

A lung nodule that remains unexplained can also create uncertainty and lead to repeated scans or additional biopsy. This follow-up is intended to confirm that the nodule is stable and that another diagnosis, including cancer, has not been missed.

Treatment can have complications as well. Antimicrobial and immune-suppressing medicines may require blood tests or other monitoring. The specific risks depend on the medicine being used.

Prevention

A necrotizing granuloma cannot always be prevented. It is a tissue response to several different problems, and some exposures are difficult to recognize or avoid.

The risk of tuberculosis can be reduced through early testing and treatment after a significant exposure. If active tuberculosis is diagnosed, following infection-control advice helps protect other people.

Soil-dwelling fungi are common in some regions. People with severely weakened immune systems may receive specific advice about avoiding heavy dust exposure, excavation sites, or activities that disturb contaminated soil. Everyday contact with soil does not mean that an infection will occur.

There is no proven general lifestyle change that prevents all necrotizing granulomas. Not smoking supports overall lung health, but smoking is not the main cause of this tissue pattern.

Living with it and what to expect

What happens over months and years is determined mainly by the cause.

An old, inactive granuloma may remain visible on scans for years without changing or causing symptoms. It may develop calcium deposits as it heals. If a solitary granuloma has been removed and no active cause is found, no further treatment may be needed after appropriate follow-up.

Treatable infections may improve with cause-specific therapy, although lung scars or nodules can remain after the infection has cleared. A persistent mark on a scan does not always mean that infection is still active.

Inflammatory conditions may have periods of improvement and recurrence. They may require longer-term monitoring of the lungs and other organs, even when symptoms are controlled.

If the initial testing is inconclusive, repeat imaging helps show whether a nodule is stable, shrinking, or growing. Keep scheduled scans and appointments even if you feel well. New symptoms between visits should be reported rather than saved for the next routine appointment.

Most day-to-day activity is guided by symptoms and the underlying diagnosis. A noninfectious granuloma is not contagious. If tuberculosis or another transmissible infection is suspected, you may receive temporary advice about masks, ventilation, or limiting close contact until testing is complete.

When to get medical help

Arrange routine medical follow-up if you have been told that a biopsy showed a necrotizing granuloma. The result needs to be interpreted alongside cultures, scans, exposure history, and other tests.

Contact a healthcare service promptly if you develop:

Seek urgent or emergency care for severe trouble breathing, a large or increasing amount of coughed-up blood, blue or grey lips, fainting, confusion, or severe chest pain.

Fever or rapidly worsening symptoms also need prompt assessment if your immune system is weakened by chemotherapy, an organ transplant, high-dose immune-suppressing treatment, or another medical condition.

- A worsening or persistent cough
- Fever, night sweats, or unexplained weight loss

- New shortness of breath
- Blood in your mucus
- New skin sores, swollen joints, dark urine, or reduced urination

Common questions

Is a necrotizing granuloma cancer?

No. It is a pattern of inflammation and tissue damage, not a type of cancer. However, a granuloma can resemble a cancerous nodule on imaging. Biopsy and follow-up may be needed to tell them apart.

Does a necrotizing granuloma mean I have tuberculosis?

Not necessarily. Tuberculosis is an important possible cause, but nontuberculous mycobacteria, fungi, inflammatory conditions, and other causes can create the same tissue pattern. Tuberculosis testing, cultures, and your exposure history help clarify the cause.

Is it contagious?

The granuloma itself is not contagious. Some underlying infections, especially active tuberculosis in the lungs or throat, can spread between people. Most fungal causes and nontuberculous mycobacteria do not usually spread from person to person.

Can a necrotizing granuloma go away?

The active inflammation may settle after the cause clears or is treated. The nodule or scar can remain visible on scans, sometimes with calcium inside it. A remaining scan abnormality does not automatically mean that treatment failed.

Why did my biopsy not find an organism?

Organisms may be sparse, absent from the sampled area, or difficult to grow. The trigger may also be noninfectious or may have already cleared. Clinicians may use cultures, molecular tests, blood tests, and follow-up imaging to investigate further.

Will I need surgery?

Not always. Surgery may be used when a nodule cannot be diagnosed safely by another method or when cancer remains a concern. Many granulomas are diagnosed with a needle biopsy or bronchoscopy and then treated or monitored without surgery.

Sources

- Mukhopadhyay S, Gal AA. "Granulomatous Lung Disease: An Approach to the Differential Diagnosis." Archives of Pathology & Laboratory Medicine . 2010;134(5):667-690.
- Ohshimo S, Guzman J, Costabel U, Bonella F. " Differential Diagnosis of Granulomatous Lung Disease: Clues and Pitfalls ." European Respiratory Review . 2017;26(145):170012.
- Lewinsohn DM, Leonard MK, LoBue PA, et al. " Official ATS/IDSA/CDC Clinical Practice Guidelines: Diagnosis of Tuberculosis in Adults and Children ." Clinical Infectious Diseases . 2017;64(2):e1-e33.
- Daley CL, Iaccarino JM, Lange C, et al. " Treatment of Nontuberculous Mycobacterial Pulmonary Disease: An Official ATS/ERS/ESCMID/IDSA Clinical Practice Guideline ." Clinical Infectious Diseases . 2020;71(4):e1-e36.
- Chung SA, Langford CA, Maz M, et al. " 2021 American College of Rheumatology/Vasculitis Foundation Guideline for the Management of ANCA-Associated Vasculitis ." Arthritis & Rheumatology . 2021;73(8):1366-1383.