CONDITION

Neuropathy secondary to drugs

Neuropathy secondary to drugs is nerve damage caused by certain medicines, often leading to numbness, tingling, pain, or weakness.

WHAT IT IS
Peripheral nerve damage caused by certain medicines

BODY SYSTEM
Nervous system

MAIN SYMPTOMS
Numbness, tingling, burning pain, weakness

WHAT CAUSES IT
Toxic effects of medicines on nerves

HOW IT IS DIAGNOSED
Clinical assessment, nerve tests, exclusion of other causes

MAIN TREATMENT
Safe medicine adjustment, symptom control, rehabilitation

CAN IT BE CURED?
Sometimes - nerves recover after medicine changes

IS IT AN EMERGENCY?
Sometimes - rapid weakness or breathing trouble

LONG-TERM OUTLOOK
Often improves, but some nerve damage persists

Overview

Neuropathy secondary to drugs is damage to peripheral nerves caused by a medicine. Peripheral nerves carry messages between your brain, spinal cord, skin, muscles, and internal organs.

The condition most often affects nerves in the feet and hands. It may cause numbness, tingling, burning pain, unusual sensitivity, or weakness. Some medicines can also affect nerves that control blood pressure, digestion, bladder function, or sweating.

Cancer treatments are a well-known cause, but some antibiotics like amoxicillin or doxycycline , antiviral medicines, heart medicines, and other drugs can also injure nerves. Risk is often related to how much medicine you receive over time, although individual susceptibility matters.

Not every new symptom during treatment is caused by the medicine. Diabetes, vitamin deficiencies, alcohol use, pressure on a nerve, infections, and other medical conditions can produce similar symptoms. A careful assessment is therefore important.

Drug-related neuropathy may improve after treatment is changed or completed. Recovery can take months, and some people have lasting symptoms.

Symptoms

Symptoms commonly begin in the toes or soles of the feet. They may later move upward or affect the fingers. Both sides are usually affected in a similar pattern, sometimes described as a "stocking and glove" distribution.

Sensory symptoms can include:

Muscle-related symptoms may include weakness, cramps, reduced grip, or difficulty lifting the front of the foot. You may find it harder to fasten buttons, open containers, climb stairs, or keep your balance.

Damage to automatic, or autonomic, nerves is less common. It can cause dizziness when standing, constipation or diarrhoea, changes in sweating, bladder problems, or sexual difficulties.

Symptoms may appear during treatment, shortly afterward, or after prolonged use. With some chemotherapy medicines, neuropathy can continue to worsen for a time after the drug has stopped. This is sometimes called "coasting."

- Numbness, tingling, or "pins and needles"
- Burning, shooting, electric, or stabbing pain
- Reduced ability to feel heat, cold, pain, or vibration
- Pain from light touch, such as bedsheets against the feet
- Poor awareness of where the feet or hands are positioned

Causes and risk factors

Drug-related neuropathy happens when a medicine or one of its breakdown products damages nerve cells, their protective covering, or the structures that transport materials along the nerve. Different medicines affect nerves in different ways.

Medicines associated with peripheral neuropathy include certain:

This is not a complete list. The likelihood and pattern of neuropathy differ greatly between medicines. A drug being associated with neuropathy does not mean everyone taking it will develop nerve damage.

Risk may be higher with a larger cumulative exposure, meaning the total amount received over time. Combining medicines that can injure nerves may also increase risk.

You may be more vulnerable if you already have neuropathy from diabetes, alcohol, a vitamin deficiency, an inherited nerve condition, or a previous treatment. Kidney or liver problems may affect how some medicines are cleared from the body. Age and genetic differences may also influence susceptibility, but they do not reliably predict who will be affected.

- Chemotherapy drugs, including platinum drugs, taxanes, vinca alkaloids, bortezomib, and thalidomide-related medicines
- Antimicrobials, such as prolonged metronidazole or linezolid treatment, nitrofurantoin, and isoniazid
- Older antiretroviral drugs used to treat HIV
- Heart medicines, including amiodarone
- Other medicines, including colchicine in some circumstances

Diagnosis

There is no single test that proves a medicine caused neuropathy. Diagnosis is based on your symptoms, neurological examination, medication history, and tests used to assess the nerves or rule out other causes.

Your clinician will usually review:

During an examination, your strength, reflexes, balance, coordination, and ability to feel touch, vibration, temperature, or pinprick may be checked.

Blood tests can look for other explanations, such as diabetes, vitamin B12 deficiency, thyroid problems, or kidney and liver disease.

Nerve conduction studies measure how quickly electrical signals travel through larger nerves. Electromyography, or EMG, assesses electrical activity in muscles. These tests can describe the type and severity of nerve damage, but they cannot always identify its cause.

Some painful neuropathies affect very small nerve fibres and may not appear on standard nerve conduction studies. In selected cases, skin biopsy or tests of autonomic nerve function may be considered.

- When the symptoms began in relation to treatment
- Which medicines you take, including non-prescription products
- The total duration or amount of the suspected medicine
- Whether symptoms changed after treatment was adjusted
- Other conditions that could affect your nerves

Treatment and management

Do not stop or reduce a prescribed medicine on your own. Suddenly stopping some medicines can be harmful, and changing an effective cancer or infection treatment requires careful consideration.

If a drug is thought to be causing neuropathy, the prescribing team may consider a treatment pause, dose reduction, schedule change, or a different medicine. The choice balances nerve symptoms against the reason the medicine is needed. Sometimes treatment must continue with close monitoring.

Pain can be treated with medicines used for nerve pain. Duloxetine has the strongest evidence for painful neuropathy caused by chemotherapy, although it is not suitable or effective for everyone. Other nerve-pain medicines or topical treatments may be considered according to your symptoms and medical history.

Ordinary painkillers often have limited effect on burning or electric nerve pain. Supplements should not be started as a substitute for assessment. Evidence for many marketed "nerve support" products is weak, and some can interact with treatment or cause neuropathy when taken in excess.

Physical therapy can help with strength, balance, walking, and safe exercise. Occupational therapy can suggest ways to manage hand symptoms and adapt daily tasks. A walking aid or brace may reduce falls when weakness or balance problems are significant.

Protect areas with reduced sensation. Check your feet regularly for blisters, cuts, pressure marks, or burns. Wear well-fitting footwear, test bath water with an unaffected area or thermometer, and use care with heaters and hot cookware.

Complications

Loss of feeling can make injuries harder to notice. Cuts, burns, or pressure sores may become infected if they are not found and treated. Balance problems and weakness can increase the

risk of falls.

Hand numbness may interfere with work, cooking, writing, or driving. Persistent pain can disturb sleep and affect concentration or mood.

Severe neuropathy can limit mobility and independence. Autonomic nerve damage may cause fainting, bowel or bladder difficulties, or problems tolerating changes in position. These effects are less common but need medical assessment.

Prevention

Drug-related neuropathy cannot always be prevented. Some necessary medicines can damage nerves even when they are used correctly.

Before starting a medicine known to affect nerves, tell the prescribing team about existing numbness, diabetes, previous chemotherapy, heavy alcohol use, nutritional problems, or an inherited nerve condition. A baseline examination can make later changes easier to identify.

Report new tingling, numbness, pain, balance problems, or weakness early. This may allow treatment to be reviewed before symptoms become more severe. Attend planned monitoring appointments and blood tests, especially if kidney or liver function affects how your medicine is processed.

No medicine or supplement has been proved to prevent all chemotherapy-induced neuropathy. Cooling or compression techniques are being studied for some chemotherapy regimens, but their use is not established for every drug or treatment centre.

Avoid taking high-dose vitamins or supplements without guidance. Excess vitamin B6, for example, can itself cause neuropathy.

Living with it and what to expect

The course depends on the medicine, the type of nerve damage, your total exposure, and whether you had neuropathy beforehand.

Some people notice improvement within weeks or months after the suspected medicine is reduced, replaced, or completed. Nerves heal slowly, so changes may be gradual. Pain may improve before normal sensation returns, or numbness may remain after pain settles.

Symptoms caused by some chemotherapy drugs can worsen for several weeks or months after treatment ends before stabilising. In other cases, neuropathy remains unchanged or persists for years. Severe nerve damage may be permanent.

Regular activity within your abilities can help preserve strength and balance, although it does not directly repair damaged nerves. Rehabilitation, practical adaptations, and good foot care can help you stay active while recovery occurs.

Keep track of symptoms that affect function, such as falls, difficulty walking, dropped objects, or problems sleeping. These details are often more useful at follow-up than a pain score alone.

When to get medical help

Arrange a routine appointment if you develop new or persistent tingling, numbness, burning pain, or reduced sensation while taking a medicine. Mention symptoms even if they are mild, particularly during chemotherapy or prolonged antimicrobial treatment.

Seek prompt medical advice if symptoms are spreading quickly, pain is becoming difficult to manage, or weakness is interfering with walking or hand use. Repeated falls, fainting, a new foot wound, or signs of infection around an unnoticed injury also need timely assessment.

Get emergency help for:

These symptoms are not typical of mild drug-related neuropathy and may have another urgent cause.

- New trouble breathing or swallowing
- Rapidly worsening weakness or inability to stand
- Sudden one-sided weakness, facial drooping, or speech difficulty
- New loss of bladder or bowel control with numbness around the groin

Common questions

Can drug-induced neuropathy go away?

Sometimes. Mild nerve injury may improve after the medicine is changed or completed. Recovery can take months because peripheral nerves heal slowly. More severe damage, especially after a large cumulative exposure, may not fully recover.

Should I stop the medicine causing my symptoms?

Do not stop it without speaking to the prescribing team. The risks of stopping depend on why you need the medicine. Your team may adjust the dose, delay treatment, switch medicines, or continue treatment while managing symptoms.

How quickly can a medicine cause neuropathy?

Timing differs by medicine. Some chemotherapy drugs can cause symptoms during or soon after an infusion, while other medicines cause neuropathy only after weeks or months of exposure. Previous nerve damage may make symptoms appear sooner.

Why is my neuropathy worse after treatment ended?

With some drugs, particularly certain platinum chemotherapy medicines, nerve injury can continue to develop after treatment stops. This "coasting" effect may last for a period before symptoms stabilise or improve.

What helps numb or painful feet?

Nerve-pain treatment may reduce burning or shooting pain, but it usually does not restore sensation. Physical therapy, suitable footwear, balance exercises, and daily skin checks can reduce the effects of numbness and help prevent falls or unnoticed injuries.

Can I drive with peripheral neuropathy?

It depends on whether you can reliably feel and operate the pedals and respond quickly. Hand weakness, numb feet, or slowed reactions may make driving unsafe. Stop driving and seek an assessment if you cannot confidently control the vehicle.

Sources

- Loprinzi CL, Lacchetti C, Bleeker J, et al. "Prevention and Management of Chemotherapy-Induced Peripheral Neuropathy in Survivors of Adult Cancers: ASCO Guideline Update." Journal of Clinical Oncology . 2020;38(28):3325-3348.
- Staff NP, Grisold A, Grisold W, Windebank AJ. "Chemotherapy-Induced Peripheral Neuropathy: A Current Review." Annals of Neurology . 2017;81(6):772-781.
- Vilholm OJ, Christensen AA, Zedan AH, Itani M. "Drug-Induced Peripheral Neuropathy." Basic & Clinical Pharmacology & Toxicology . 2014;115(2):185-192.
- National Cancer Institute. "Peripheral Neuropathy and Cancer Treatment." Patient information on cancer treatment side effects.
- National Institute of Neurological Disorders and Stroke. "Peripheral Neuropathy." Overview of symptoms, diagnosis, treatment, and nerve recovery.