CONDITION

Norrbottnian gaucher disease

Norrbottnian gaucher disease is an inherited form of Gaucher type 3 that affects the organs, bones, blood, and nervous system.

WHAT IT IS
Inherited chronic neuronopathic Gaucher condition

BODY SYSTEM
Nervous system, blood, bones, liver, and spleen

MAIN SYMPTOMS
Enlarged organs, abnormal eye movements, bone pain, seizures

WHAT CAUSES IT
Inherited GBA1 gene changes

HOW IT IS DIAGNOSED
Enzyme activity testing and GBA1 genetic testing

MAIN TREATMENT
Enzyme replacement and supportive care

CAN IT BE CURED?
No - treatment controls but does not remove it

IS IT AN EMERGENCY?
Sometimes - prolonged seizure, severe bleeding, or breathing trouble

LONG-TERM OUTLOOK
Lifelong; neurological changes may progress despite treatment

Overview

Norrbottnian gaucher disease is a form of Gaucher type 3, also called chronic neuronopathic Gaucher disease. "Neuronopathic" means that it affects the brain and nervous system as well as other parts of the body.

The condition was first described in families from Norrbotten in northern Sweden. It is related to a genetic change that became more common in that population through a founder effect. A founder effect happens when a gene variant is passed down within a relatively small or historically isolated population. The same or similar form can occur in people without Swedish ancestry.

Gaucher conditions affect an enzyme called glucocerebrosidase. Enzymes are proteins that help cells carry out chemical processes. When glucocerebrosidase does not work well, fatty

substances build up inside certain immune cells. These enlarged cells are called Gaucher cells.

Gaucher cells can collect in the spleen, liver, bone marrow, and bones. In type 3, the nervous system is also affected. This makes Norrbottnian Gaucher more complex than Gaucher type 1, which does not usually cause primary neurological symptoms.

It is a lifelong condition. Treatment can greatly improve enlargement of the liver and spleen, low blood counts, and some bone problems. Current enzyme treatment does not reach the brain in meaningful amounts, so it may not stop neurological changes.

Symptoms

Symptoms often begin during childhood, but their timing and severity are not identical in every person. Some problems develop slowly and may be noticed only during specialist examinations.

A characteristic early feature is difficulty moving the eyes quickly from side to side. These rapid eye movements are called saccades. A child may turn their head instead of shifting their eyes, or may have trouble following fast-moving objects. Up-and-down eye movements can also become affected.

Other neurological symptoms may include:

The spleen and liver may become enlarged. An enlarged spleen can lower the number of blood cells by trapping or removing them too quickly. Low red blood cells can cause tiredness, weakness, or shortness of breath. Low platelets can lead to easy bruising, nosebleeds, or bleeding that takes longer to stop.

Bone involvement can cause deep bone pain, reduced bone strength, fractures, or damage to a joint. Some people have severe episodes of bone pain called bone crises. Growth and puberty may be delayed in children with poorly controlled disease.

Not everyone develops every symptom. The neurological course can be slow in some people and more limiting in others.

- Poor balance, clumsiness, or an unsteady walk
- Muscle stiffness or unusual muscle movements
- Tremor or brief muscle jerks called myoclonus
- Seizures
- Speech or swallowing difficulties
- Changes in learning, attention, or thinking

Causes and risk factors

Norrbottnian Gaucher is caused by disease-causing changes in both copies of the GBA1 gene. This gene provides instructions for making glucocerebrosidase.

The condition follows an autosomal recessive inheritance pattern. This means a person usually inherits one altered gene copy from each parent. Parents who each carry one altered copy generally do not have Gaucher symptoms.

When both parents are carriers, each pregnancy has:

These chances start again with every pregnancy. They do not change according to the sex of the child or what happened in an earlier pregnancy.

A particular GBA1 variant historically associated with the Norrbottnian form is often called L444P in older medical reports. It may be written as p.Leu483Pro under current naming systems. Genetic reports need careful interpretation because the GBA1 gene has a nearby, highly similar pseudogene that can complicate testing.

Lifestyle, diet, parenting, infection, and environmental exposure do not cause this inherited condition.

- A 1 in 4 chance of a child inheriting the condition
- A 1 in 2 chance of a child being an unaffected carrier
- A 1 in 4 chance of a child inheriting neither altered copy

Diagnosis

Diagnosis usually combines symptoms, examination findings, enzyme testing, and genetic testing. An eye movement examination is especially important when type 3 Gaucher is suspected.

The diagnosis is confirmed by showing low glucocerebrosidase activity in white blood cells or another validated cell sample. Molecular genetic testing is then used to identify disease-causing variants in both copies of GBA1 . A screening result from a dried blood spot may need confirmation using a more specific sample and laboratory method.

Blood tests may show anaemia, low platelets, or signs of liver involvement. Biomarkers such as glucosylsphingosine can help support the diagnosis and monitor the condition, but they do not replace enzyme and genetic confirmation.

Further assessment may include:

Testing relatives may identify other affected family members or carriers. Genetic counselling can explain inheritance, testing choices, and what a result may mean for the wider family.

- MRI scans to measure the liver and spleen or assess bones
- Bone density testing
- Neurological, developmental, hearing, and eye movement assessments
- An electroencephalogram, or EEG, if seizures are suspected
- Heart and lung tests when symptoms or the person's subtype make these relevant

Treatment and management

Care is usually coordinated by a specialist metabolic team. Neurology, haematology, bone, eye, rehabilitation, and genetic services may also be involved.

Enzyme replacement therapy, or ERT, supplies a manufactured form of the missing enzyme through regular intravenous infusions. It can reduce liver and spleen enlargement, improve blood counts, and lower the risk of some bone complications. Response is monitored through symptoms, growth, blood tests, organ measurements, and bone assessments.

ERT does not cross the blood-brain barrier in useful amounts. The blood-brain barrier is a protective boundary between the bloodstream and brain. As a result, ERT does not reliably prevent or reverse abnormal eye movements, seizures, or other neurological features.

Supportive treatment is matched to the problems present. It may include anti-seizure medicines, pain management, physical or occupational therapy, speech and swallowing support, and nutritional help. Mobility aids or educational support can help a person remain active and independent.

Oral substrate reduction medicines decrease production of the fatty substance that accumulates in Gaucher cells. Available medicines have specific approvals and limitations. They are not established treatments for the neurological features of type 3 and are not interchangeable with ERT.

Bone marrow or stem cell transplantation was used historically in some people with the Norrbottnian form. It carries substantial risks and is now rarely considered for Gaucher treatment. It also cannot be expected to reverse established neurological damage.

Gene-based and brain-targeted treatments are under study, but they are not currently routine cures.

Complications

Possible complications reflect the condition's effects on the nervous system, blood, bones, and internal organs.

Neurological problems may gradually affect walking, coordination, communication, swallowing, or day-to-day activities. Seizures can become difficult to control in some people. Swallowing problems may allow food or liquid to enter the lungs, leading to aspiration and chest infections.

Long-term bone complications can include reduced bone density, fractures, chronic pain, and osteonecrosis. Osteonecrosis means that part of a bone loses its blood supply and becomes damaged, often near a major joint.

A very enlarged spleen can worsen low blood counts and cause abdominal discomfort. Severe bleeding or rupture of the spleen is uncommon but requires urgent care. Removing the spleen is generally avoided unless there is a compelling reason, because it increases infection risk and may contribute to other Gaucher complications.

Liver scarring and lung or heart complications can occur, although they are not present in everyone. Regular monitoring aims to find treatable changes before they cause major symptoms.

Prevention

Norrbottnian Gaucher cannot be prevented through lifestyle changes because it is inherited.

Carrier testing and genetic counselling can help people understand the chance of passing on a GBA1 variant. Reproductive options may include testing during pregnancy or testing embryos created through in vitro fertilisation. These are personal choices, and availability differs by country.

Early diagnosis and treatment cannot remove the genetic cause, but they may prevent or reduce some blood, organ, and bone complications.

Living with it and what to expect

Norrbottnian Gaucher needs lifelong specialist follow-up. Monitoring is used to assess blood counts, liver and spleen size, growth, nutrition, bone health, movement, eye function, learning, seizures, and swallowing.

The blood and organ features often improve with ERT. Bone recovery can take longer, and

existing joint or bone damage may not fully reverse. Neurological changes may continue even when blood tests and organ size have improved.

Some people have slowly progressive neurological symptoms and live well into adulthood. Others need increasing support with mobility, communication, seizures, or daily activities. The course cannot be predicted precisely from the name of the condition or from one genetic result.

School or work adjustments may be useful for fatigue, infusions, movement difficulties, or learning needs. Infusion schedules can also affect travel and family routines. Home infusion may be available in some health systems after a person is medically stable and appropriately assessed.

Routine appointments remain important even when symptoms seem unchanged. New seizures, increasing falls, coughing during meals, worsening pain, or reduced stamina should be discussed promptly rather than waiting for the next scheduled review.

When to get medical help

Arrange a routine or prompt medical review for new bruising, increasing tiredness, bone pain, falls, changes in walking, new learning difficulties, or possible seizure activity. Coughing or choking with food and drink also needs assessment because it may signal a swallowing problem.

Seek urgent medical help for:

Call emergency services for a seizure lasting five minutes or longer, repeated seizures without recovery, severe difficulty breathing, loss of consciousness, major bleeding, or sudden severe abdominal pain with faintness. Follow an existing emergency seizure plan if one has been provided.

- A first seizure or a clear change in a known seizure pattern
- Bleeding that does not stop with firm pressure
- Sudden severe bone or joint pain
- Repeated choking, fever after aspiration, or worsening breathing symptoms
- Severe abdominal pain, especially after an injury

Common questions

Is Norrbottnian Gaucher the same as Gaucher type 3?

It is a historically recognised form of Gaucher type 3, first described in northern Sweden. Type 3 includes other genetic and clinical patterns, so the terms are related but not always identical.

Can Norrbottnian Gaucher be cured?

There is currently no established cure. ERT can control many problems involving the blood, spleen, liver, and bones, but it does not correct the inherited gene change or reliably treat the brain.

Does it always affect the brain?

Norrbottnian Gaucher is classed as neuronopathic because neurological involvement is part of the condition. Abnormal eye movements may be the earliest or most noticeable sign. The extent and speed of other neurological changes differ between people.

Can my children inherit it?

A child must usually inherit an altered GBA1 copy from both parents to have the condition. If you have Norrbottnian Gaucher, your children will inherit at least one altered copy, but their other parent's carrier status affects whether they could inherit the condition.

How is it different from Gaucher type 1?

Both forms can enlarge the liver and spleen and affect blood counts and bones. Type 3 also causes primary neurological features, especially abnormal eye movements and, in some people, seizures or movement problems.

Will enzyme replacement stop the condition getting worse?

ERT often improves or stabilises organ, blood, growth, and bone features. It does not enter the brain effectively, so neurological symptoms may still progress. Regular neurological follow-up remains necessary even when the rest of the condition is well controlled.

Sources

- Hughes DA, Pastores GM. "Gaucher Disease." GeneReviews® . University of Washington, Seattle; updated December 2023.
- Dreborg S, Erikson A, Hagberg B. "Gaucher disease-Norrbottnian type. I. General clinical description." European Journal of Pediatrics . 1980;133:107-118.
- Stirnemann J, Belmatoug N, Camou F, et al. "A Review of Gaucher Disease Pathophysiology, Clinical Presentation and Treatments." International Journal of Molecular Sciences . 2017;18(2):441.



- Vellodi A, Tylki-Szyma□ ska A, Davies EH, et al. "Management of neuronopathic Gaucher disease: revised recommendations." Journal of Inherited Metabolic Disease . 2009;32:660-664.
- Kaplan P, Baris H, De Meirleir L, et al. "Revised recommendations for the management of Gaucher disease in children." European Journal of Pediatrics . 2013;172:447-458.
- European Medicines Agency. "Cerezyme: EPAR product information." Imiglucerase prescribing and safety information.