CONDITION

Pornography addiction

Pornography addiction describes persistent difficulty controlling pornography use despite harm to daily life, relationships, or wellbeing.

WHAT IT IS
Persistent loss of control over pornography use

BODY SYSTEM
Brain, behaviour, relationships, and sexual health

MAIN SYMPTOMS
Loss of control, failed cutbacks, life disruption, continued use

WHAT CAUSES IT
Interacting psychological, social, and learned habit factors

HOW IT IS DIAGNOSED
Clinical assessment of control, distress, impairment, and other causes

MAIN TREATMENT
Psychological therapy, behaviour changes, and related-condition care

CAN IT BE CURED?
Sometimes - lasting control and recovery are possible

IS IT AN EMERGENCY?
Sometimes - danger of self-harm or non-consensual sexual behaviour

LONG-TERM OUTLOOK
Often manageable, though urges and setbacks may recur

Overview

Pornography addiction is a commonly used term for pornography use that feels out of control and continues despite serious consequences. It can affect relationships, work, study, sleep, finances, sexual wellbeing, or mental health.

"Pornography addiction" is not a specific diagnosis in the main psychiatric classification used in the United States, the DSM-5-TR. The World Health Organization recognizes compulsive sexual behaviour disorder , which may include compulsive pornography use. It classifies this as an impulse-control disorder rather than an addiction.

This distinction matters because frequent pornography use alone does not establish a condition. Neither does having a strong sex drive, masturbating, or feeling uncomfortable about pornography for religious, cultural, or personal reasons.

The central issue is an ongoing loss of control combined with meaningful harm or disruption. Distress caused only by moral disapproval is not enough to diagnose compulsive sexual behaviour disorder. However, that distress can still be real and may deserve sensitive support.

The word "addiction" can help some people describe their experience. Others find terms such as problematic pornography use or compulsive pornography use more accurate and less stigmatizing.

Symptoms

The pattern usually involves more than simply spending a lot of time viewing pornography. A person may repeatedly decide to reduce or stop, then find that they cannot follow through.

Possible signs include:

Some people continue even when viewing pornography brings little satisfaction. They may feel brief relief followed by shame, frustration, sadness, or anxiety.

A high level of sexual interest is not, by itself, a symptom. The amount or type of pornography viewed also cannot confirm a condition without looking at control, circumstances, distress, and effects on daily life.

- Strong urges that are difficult to redirect
- Repeated unsuccessful attempts to cut back
- Pornography becoming a major focus of daily life
- Continued use despite relationship, work, financial, or emotional harm
- Neglecting sleep, responsibilities, hobbies, or social contact
- Using for longer than intended or in situations that carry serious risks
- Feeling that the behaviour has become automatic rather than chosen

Causes and risk factors

There is no single proven cause. Compulsive use may develop through an interaction between learned habits, emotional coping, sexual arousal, easy access, and a person's wider mental and social circumstances.

Pornography can provide rapid distraction or relief from boredom, loneliness, stress, anxiety, or difficult emotions. Repeating that response may strengthen a habit: a trigger leads to an urge, viewing brings short-term relief, and the same response becomes more likely next time.

Factors that may contribute include:

Certain health conditions can cause increased sexual drive or reduced inhibition. Examples include a manic episode, some neurological conditions, and effects from certain medicines or substances. These require assessment because treatment may differ.

Personal values also shape how pornography use is experienced. Conflict between behaviour and moral or religious beliefs can produce intense shame even when there is no broader loss of control. In other cases, moral conflict and compulsive behaviour may exist together.

- Depression, anxiety, trauma-related symptoms, or loneliness
- Attention-deficit/hyperactivity disorder or difficulty controlling impulses
- Relationship conflict or sexual dissatisfaction
- Privacy, constant internet access, and repeated exposure to sexual material
- Using pornography to avoid distress rather than address it
- Alcohol or drug use that reduces judgment or self-control

Diagnosis

There is no blood test, brain scan, or online questionnaire that can confirm pornography addiction.

Assessment is based on a private conversation about what is happening, how long it has continued, and how it affects daily life. A qualified health professional may ask about control over the behaviour, unsuccessful attempts to change it, triggers, consequences, sexual health, relationships, mood, sleep, medicines, and substance use.

For compulsive sexual behaviour disorder, clinicians look for a persistent failure to control repeated sexual urges or behaviours over an extended period, generally six months or longer. The pattern must cause significant distress or impair important areas of life. Distress based only on moral judgment is not sufficient.

Assessment also helps distinguish the problem from other conditions. Increased sexual activity may occur during mania, for example. Repetitive sexual thoughts may sometimes be related to obsessive-compulsive disorder. Substance use, trauma, depression, anxiety, or medication effects can also influence the pattern.

Screening questionnaires may organize information, but they should not be used alone to label someone. Pornography use is personal, and a respectful assessment should avoid assumptions about sexuality, relationships, culture, or beliefs.

Treatment and management

Treatment aims to restore choice and reduce harm. Depending on the person's goals, this may mean stopping pornography use or reducing it to a level that no longer disrupts life.

Psychological therapy is the main treatment approach. Cognitive behavioural therapy, often called CBT, can help identify triggers, challenge unhelpful thinking, manage urges, and develop alternative responses. Acceptance-based approaches may help a person experience an urge without automatically acting on it.

Treatment may also address shame, loneliness, trauma, relationship difficulties, or sexual concerns. If depression, anxiety, ADHD, bipolar disorder, substance use, or another condition is contributing, treating that condition can be an important part of recovery.

Practical changes may support therapy. These can include:

Blocking software can create useful time to reconsider an urge, but it rarely addresses the emotional or behavioural pattern by itself.

Couples or psychosexual therapy may be helpful when the problem has affected trust, intimacy, or communication. A partner should not be expected to monitor another adult's devices or take responsibility for controlling their behaviour.

No medicine is specifically approved as a cure for pornography addiction. Specialists sometimes consider medication for a related mental health condition or, in selected cases, to reduce compulsive sexual symptoms. The choice requires individual assessment because benefits and risks differ.

Peer-support groups help some people feel less isolated and provide structure. Their methods and quality differ, so they are best viewed as possible support rather than a replacement for professional care.

- Tracking triggers and high-risk situations without self-criticism
- Changing routines linked to viewing
- Keeping devices out of bedrooms or other private settings
- Using filters, limits, or blocking tools as added barriers
- Planning another activity for times when urges are strongest
- Rebuilding sleep, exercise, social contact, and valued hobbies

Complications

Persistent compulsive pornography use can consume time and attention. It may contribute to missed work, reduced study performance, sleep loss, financial spending, secrecy, and withdrawal from other activities.

Relationships may be affected by broken agreements, reduced trust, dishonesty, differences in sexual expectations, or less interest in shared intimacy. These effects are not inevitable, and pornography use does not affect every relationship in the same way.

Some people report erection difficulties, reduced arousal with a partner, delayed orgasm, or pressure to recreate what they have viewed. Research has not shown that pornography is the sole cause of sexual difficulties in every case. Anxiety, physical health, medicines, relationship factors, and other sexual habits may also contribute.

Shame can make the pattern worse by increasing isolation and discouraging help-seeking. Severe distress may contribute to depression or thoughts of self-harm.

Viewing illegal sexual material can cause serious harm to victims and have major legal consequences. Any concern about losing control around non-consensual behaviour or sexual material involving children requires immediate action to protect others.

Prevention

There is no proven way to prevent every case of compulsive pornography use. It is not simply caused by weak willpower, and it is not an inherited disease with a specific preventive test.

Risk may be reduced by noticing early when pornography becomes the main way of coping with stress, loneliness, boredom, or difficult feelings. Keeping a balanced routine, setting personal boundaries around device use, and seeking help for mental health or relationship problems may prevent a habit from becoming more disruptive.

Age-appropriate education about consent, realistic sexual expectations, online privacy, and the difference between pornography and real-life intimacy can support healthier decisions. Filters and parental controls can limit children's exposure, but open, non-shaming conversations are also important.

Living with it and what to expect

Change often takes place over months rather than days. Early progress may involve understanding triggers, making access less automatic, and learning to tolerate urges without acting on them. Urges may still occur even when behaviour is becoming more controlled.

A setback does not erase progress. It can show where a plan needs adjustment, such as during stress, isolation, alcohol use, travel, or conflict. Repeated setbacks may mean that more structured professional support is needed.

Over time, many people develop greater control and spend less mental energy on pornography. Sleep, concentration, relationships, and sexual confidence may improve when they have been affected. The pace and extent of improvement are not the same for everyone.

Some people choose long-term abstinence. Others work toward non-compulsive use that fits their values and relationship agreements. A useful goal is not a particular number of days or hours, but a stable return of choice, safety, and everyday functioning.

Shame and secrecy can make recovery harder. Support that respects sexuality while addressing harmful behaviour is more likely to encourage honest, lasting change.

When to get medical help

Arrange a routine appointment with a primary care or mental health professional if pornography use repeatedly feels beyond your control, causes significant distress, or interferes with work, study, sleep, finances, relationships, or sexual wellbeing.

Routine help is also appropriate if attempts to stop lead to repeated cycles of secrecy and shame, or if depression, anxiety, trauma symptoms, substance use, or attention problems may be contributing.

Seek urgent mental health help if distress is becoming overwhelming, you are having thoughts of self-harm, or your behaviour is escalating into unsafe situations.

Get emergency help now if you may act on thoughts of suicide, cannot keep yourself safe, or believe you may engage in sexual behaviour without consent or involving a child. Move away from potential victims, avoid being alone with anyone at risk, and contact local emergency services or a crisis service.

Common questions

Am I addicted if I watch pornography every day?

Not necessarily. Frequency alone does not determine whether there is a condition. The more important questions are whether you can choose not to view it and whether it is causing significant harm or disruption.

Is pornography addiction an official diagnosis?

Not under that exact name in major diagnostic systems. The World Health Organization recognizes compulsive sexual behaviour disorder, which can include problematic pornography use. It is classified as an impulse-control disorder, not an addiction.

Can pornography cause erectile dysfunction?

Some people with problematic use report erection or arousal difficulties, especially with a partner. Pornography is not proven to be the only cause. Physical health, anxiety, medicines, relationship concerns, and other factors should also be considered.

Do I have to stop completely?

Not everyone chooses the same goal. Some people find abstinence clearer and easier to maintain. Others aim for controlled use that does not interfere with life or conflict with relationship agreements. Treatment should focus on safety, control, and your informed goals.

Are irritability and strong urges withdrawal symptoms?

People who reduce frequent use may experience urges, restlessness, irritability, low mood, or difficulty concentrating. These experiences can be uncomfortable, but pornography withdrawal is not defined in the same way as withdrawal from alcohol or certain drugs.

Can a relationship recover after secret pornography use?

Recovery is possible, but rebuilding trust may take time. Honest communication, clear agreements, accountability, and couples or psychosexual therapy can help. No partner is required to remain in a relationship or accept boundaries that feel unsafe.

Sources

- World Health Organization. ICD-11 for Mortality and Morbidity Statistics: Compulsive sexual behaviour disorder, code 6C72
- World Health Organization. Clinical Descriptions and Diagnostic Requirements for ICD-11 Mental, Behavioural and Neurodevelopmental Disorders . 2024
- Kraus SW, Krueger RB, Briken P, and colleagues. "Compulsive sexual behaviour disorder in the ICD-11." World Psychiatry . 2018;17(1):109-110
- Antons S, Engel J, Briken P, Krüger THC, Brand M, and Stark R. "Compulsive sexual behaviour disorder." Nature Reviews Disease Primers . 2022;8:70
- Grubbs JB, Perry SL, Wilt JA, and Reid RC. "Pornography problems due to moral incongruence: An integrative model with a systematic review and meta-analysis." Archives of Sexual Behavior . 2019;48:397-415

- American Association of Sexuality Educators, Counselors and Therapists. Position Statement on Sex Addiction . 2016
- Hallberg J, Kaldo V, Arver S, Dhejne C, and Öberg KG. "A randomized controlled study of group-administered cognitive behavioral therapy for hypersexual disorder in men." The Journal of Sexual Medicine . 2019;16(5):733-745