CONDITION

Presenile tremor syndrome

Presenile tremor syndrome is an older label for persistent shaking that begins before older age, usually during movement.

WHAT IT IS
Persistent action tremor beginning before older age

BODY SYSTEM
Nervous system

MAIN SYMPTOMS
Shaky hands, action tremor, head tremor, voice tremor

WHAT CAUSES IT
Often unknown, sometimes inherited

HOW IT IS DIAGNOSED
Neurological examination and exclusion of other causes

MAIN TREATMENT
Practical strategies, medicines, and sometimes procedures

CAN IT BE CURED?
No - treatment can reduce symptoms

IS IT AN EMERGENCY?
Sometimes - sudden tremor with stroke symptoms

LONG-TERM OUTLOOK
Often slowly progressive without shortening life

Overview

Presenile tremor syndrome is an older and imprecise term for ongoing shaking that begins before older age. "Presenile" means occurring earlier than expected in later life. There is no single accepted age cutoff.

The term is not widely used in current movement-disorder classifications. In many records, it describes a pattern now assessed as essential tremor : shaking that is most noticeable when holding a position or making a movement. However, the old label does not confirm a specific modern diagnosis.

A tremor is an involuntary, rhythmic movement of part of the body. It can affect the hands, arms, head, voice, jaw, or legs. With this type of tremor, shaking is often more obvious while writing, eating, drinking, using tools, or holding the arms out.

The condition usually does not shorten life. It matters because tremor can interfere with daily tasks, work, hobbies, and confidence. It can also resemble shaking caused by medicines, thyroid problems, Parkinson's disease, or other neurological conditions. A careful assessment helps clarify the cause.

Symptoms

The main symptom is rhythmic shaking. It may be mild enough to notice only during precise tasks, or strong enough to spill drinks and make handwriting difficult.

Typical features may include:

Tremor may become more noticeable with stress, tiredness, hunger, caffeine, illness, or strong emotion. This does not mean anxiety caused the underlying condition. These factors can temporarily increase a tremor that is already present.

A person may have periods when symptoms are less troublesome. Tremor often becomes more noticeable over years, but the pace is not predictable. One side can be worse even when both sides are affected.

Shaking mainly at rest, marked stiffness, slowed movement, weakness, loss of coordination, or a major change in walking is less typical of uncomplicated essential tremor. These features do not identify the cause by themselves, but they are reasons for another neurological assessment.

- Shaking of both hands or arms during movement
- Tremor while holding the arms or an object in one position
- Head shaking, such as a "yes-yes" or "no-no" movement
- A wavering or shaky voice

Causes and risk factors

The exact cause is often unknown. Current research suggests that essential-type tremor involves altered signalling within brain networks that coordinate movement. It is not caused by weak muscles or a lack of willpower.

Tremor can run in families. A person may inherit a tendency to develop it, although the genes involved are not fully understood. Having a family history does not predict exactly when symptoms will start or how troublesome they will become.

The older label "presenile tremor syndrome" can also hide other causes of shaking. These include:

Some medicines merely make an existing tremor more visible. Do not stop a prescribed medicine suddenly unless a healthcare professional advises you to do so.

- Medicines such as some stimulants like methylphenidate or lisdexamfetamine , asthma treatments, antidepressants like sertraline or fluoxetine , or mood stabilisers
- An overactive thyroid gland
- Low blood sugar or other metabolic problems
- Excess caffeine or nicotine
- Alcohol or sedative withdrawal
- Parkinsonian, dystonic, or coordination-related tremors

Diagnosis

There is no blood test or scan that confirms presenile tremor syndrome or essential tremor. Diagnosis is based mainly on the history of the shaking and a neurological examination.

The clinician may ask when the tremor began, which body parts are affected, and whether it occurs at rest, while holding a posture, or during movement. They may watch you write, draw a spiral, pour water, hold out your arms, or bring a finger toward a target.

The examination also checks muscle tone, movement speed, coordination, reflexes, walking, and balance. This helps identify signs that point toward another type of tremor.

Under current specialist classifications, essential tremor usually means an action tremor affecting both upper limbs for at least three years, without other neurological signs that better explain it. A shorter history can still be assessed and treated, but the final classification may remain uncertain while symptoms are followed.

Tests are selected according to the individual situation. Blood tests may check thyroid function or other possible metabolic causes. Brain imaging may be considered when tremor began suddenly, followed an injury, or occurs with unusual examination findings. A specialised dopamine-transporter scan is occasionally used when it is difficult to distinguish essential tremor from a parkinsonian condition, but it does not confirm essential tremor itself.

Because the term is outdated, someone who finds it in an old medical record may benefit from having the diagnosis reviewed using current terminology.

Treatment and management

Treatment is guided by how much the tremor affects daily life. Mild tremor may not need medical treatment. Monitoring and practical adjustments can be enough.

Useful changes can include using cups with lids, heavier utensils, wide-grip pens, voice controls, or two hands for tasks. Resting the elbows on a surface may improve control. Occupational therapy can suggest equipment and techniques suited to work, cooking, dressing, or writing.

Reducing caffeine may help if it clearly worsens symptoms. Regular meals, sleep, and planning demanding tasks for steadier times can also make tremor easier to manage.

Medicines commonly used for essential-type tremor include certain beta blockers and anti-seizure medicines. Other medicines may be considered when these do not help or are not suitable. Benefits differ between people, and side effects can limit treatment. A clinician should review other health conditions and medicines before recommending an option.

Botulinum toxin injections may help selected head, voice, or hand tremors. The injections temporarily weaken specific overactive muscles, so unwanted weakness can occur.

For severe tremor that remains disabling despite medicines, specialist procedures may be considered. Deep brain stimulation uses implanted electrodes to adjust movement-related brain signals. Focused ultrasound creates a small, targeted treatment area in the brain without an implanted device. These procedures have important benefits, limitations, and risks and require assessment by an experienced movement-disorder team.

Alcohol may briefly reduce some essential tremors, but it is not a safe treatment. Tremor can rebound as alcohol wears off, and regular use can lead to dependence and other health problems.

Complications

The most common complications come from loss of fine control. Tremor may make eating, drinking, shaving, applying makeup, fastening clothes, typing, or using tools harder.

Spills, cuts, and burns can occur when handling hot liquids or sharp objects. Practical adaptations can reduce these risks without requiring someone to stop valued activities.

Visible shaking may cause embarrassment or lead someone to avoid meals, meetings, or social events. Support for anxiety or low mood can be helpful when these become part of the experience.

Treatment can also cause complications. Medicines may produce tiredness, dizziness, slowed thinking, or other effects, depending on the drug. Injections and procedures carry their own risks. These should be weighed against how much the tremor affects everyday life.

A new problem with balance, memory, speech, stiffness, or movement should not automatically be blamed on the existing tremor. It deserves reassessment.

Prevention

A primary or inherited tremor cannot currently be prevented. There is no proven diet, supplement, or exercise programme that stops it from developing.

You may be able to reduce temporary worsening by identifying personal triggers such as excess caffeine, poor sleep, skipped meals, or stress. This manages symptoms rather than preventing the condition.

Medication reviews can sometimes identify a drug that is increasing tremor. Any change should be planned with the clinician who prescribed it, especially when stopping suddenly could cause withdrawal or a return of the condition being treated.

Living with it and what to expect

Presenile or essential-type tremor is usually long-lasting. Symptoms may remain mild for years or gradually become more noticeable. Progression is often slow, but no individual course can be promised.

The tremor may spread from the hands to the head or voice over time. It does not necessarily affect every body part. Strength and sensation usually remain intact, although precise movements can become harder because of the shaking.

Many people continue working, driving, exercising, and living independently. Whether an activity remains safe depends on the severity of the tremor and the task involved. Equipment changes, occupational therapy, and treatment adjustments can preserve independence.

Follow-up is useful when symptoms change or treatment is being used. A routine review may cover daily function, side effects, new medicines, and whether the tremor still fits its original classification. Treatment can be increased, reduced, or changed as needs evolve.

When to get medical help

Arrange a routine medical appointment if a tremor:

Seek urgent medical advice if a new tremor develops soon after starting or increasing a medicine and is accompanied by fever, marked agitation, muscle rigidity, vomiting, or a very fast heartbeat.

Call emergency services for sudden shaking accompanied by facial drooping, one-sided weakness or numbness, trouble speaking, severe loss of balance, confusion, seizure, or a sudden severe headache. These symptoms may indicate a stroke or another emergency rather than a chronic tremor condition.

- Persists, worsens, or begins affecting daily activities
- Starts after a medicine change
- Interferes with eating, drinking, writing, work, or sleep
- Occurs with stiffness, slowing, balance problems, or unusual movements
- Has not previously been assessed

Common questions

Is presenile tremor syndrome the same as essential tremor?

Not always, because presenile tremor syndrome is an older, non-specific label. In many records it appears to describe an essential-type action tremor beginning before older age. A current assessment can determine which modern tremor classification fits best.

Is this an early sign of Parkinson's disease?

Essential-type tremor is different from the typical tremor of Parkinson's disease. Essential tremor is usually most obvious during posture or movement, while Parkinsonian tremor often appears at rest and may occur with slowness or stiffness. These patterns can overlap, so the type cannot be confirmed from one symptom alone.

Will the shaking keep getting worse?

It may gradually worsen, but the rate differs. Some people have mild, stable symptoms for long periods, while others need more support over time. A rapid or sudden change is not typical and should be assessed.

Can stress cause this tremor?

Stress can make shaking more visible, but it does not usually explain a persistent essential-type tremor by itself. Tiredness, caffeine, illness, and strong emotions can have a similar temporary effect.

Can exercise make tremor better?

Exercise does not cure the condition, but it can support strength, balance, sleep, and general health. Tremor may briefly increase after strenuous activity. A physiotherapist can help if shaking or balance problems make exercise difficult.

Can I still drive with a tremor?

Many people can drive safely when tremor is mild and control of the vehicle is not affected. Driving should be reconsidered if shaking interferes with steering, operating controls, reacting quickly, or taking medicines without impairing side effects. Local licensing rules may also apply.

Sources

- Bhatia KP, Bain P, Bajaj N, et al. Consensus Statement on the classification of tremors from the International Parkinson and Movement Disorder Society . Movement Disorders . 2018;33(1):75-87.
- Ferreira JJ, Mestre TA, Lyons KE, et al. MDS evidence-based review of treatments for essential tremor . Movement Disorders . 2019;34(7):950-958.
- Zesiewicz TA, Elble RJ, Louis ED, et al. Evidence-based guideline update: Treatment of essential tremor . Neurology . 2011;77(19):1752-1755.
- National Institute of Neurological Disorders and Stroke. Tremor . US National Institutes of Health.
- NHS. Tremor or shaking hands . National Health Service.