CONDITION

Schizoaffective disorder

Schizoaffective disorder is a long-term mental health condition involving psychosis alongside major episodes of depression or mania.

WHAT IT IS
Long-term condition combining psychosis with major mood episodes

BODY SYSTEM
Mental health and nervous system

MAIN SYMPTOMS
Hallucinations, delusions, depression or mania, disorganised thinking

WHAT CAUSES IT
Unknown; genetic, brain, and environmental factors contribute

HOW IT IS DIAGNOSED
Psychiatric assessment over time; tests exclude other causes

MAIN TREATMENT
Antipsychotic medicine, mood treatment, therapy, and practical support

CAN IT BE CURED?
No - symptoms can often be managed

IS IT AN EMERGENCY?
Sometimes - suicidal intent or immediate danger

LONG-TERM OUTLOOK
Often long-term, with episodes, recovery, and ongoing support

Overview

Schizoaffective disorder is a mental health condition with symptoms of both psychosis and a major mood disorder.

Psychosis affects how you interpret reality. It can include hallucinations, such as hearing voices, or delusions, which are strongly held beliefs that other people do not share. Mood episodes involve either depression, mania, or both.

Mania is a period of unusually high or irritable mood, increased energy, reduced need for sleep, and changes in behaviour. Depression can cause persistent low mood, loss of interest, low energy, and thoughts of death or suicide.

In schizoaffective disorder, psychosis also occurs for a period when a major mood episode is

not present. This timing helps distinguish the condition from depression or bipolar disorder with psychotic features.

There are two main types:

The condition can affect relationships, work, education, physical health, and daily tasks. Treatment can reduce symptoms and help many people build a stable and meaningful life.

- Bipolar type includes manic episodes and may also include depression
- Depressive type includes major depressive episodes but no history of mania

Symptoms

Symptoms differ between people and can change over time. They may develop gradually or become noticeable during an acute episode.

Psychotic symptoms can include hallucinations. Hearing voices is the most common example, but a person may also see, feel, smell, or taste something that others do not.

Delusions may involve feeling watched, followed, threatened, or controlled. A person may believe that messages have a special personal meaning. These experiences can feel completely real.

Disorganised thinking may make speech difficult to follow. Concentration, planning, memory, or decision-making may also become harder. Behaviour can seem confused or poorly matched to the situation.

Depressive symptoms may include:

Manic symptoms may include:

Some people also experience reduced motivation, limited emotional expression, social withdrawal, or difficulty enjoying activities. These are sometimes called negative symptoms because ordinary abilities or responses have become less noticeable.

- Persistent sadness, emptiness, or irritability
- Loss of interest or pleasure
- Low energy and changes in sleep or appetite
- Feelings of guilt, worthlessness, or hopelessness
- Thoughts of self-harm or suicide
- Very high, excited, or irritable mood
- Much more energy or activity than usual

- Needing little sleep without feeling tired
- Rapid speech, racing thoughts, or easy distraction
- Impulsive spending, sexual behaviour, travel, or risky decisions

Causes and risk factors

The exact cause is unknown. Schizoaffective disorder probably develops through a combination of genetic, brain, developmental, and environmental factors.

It can run in families, but it is not inherited in a simple or predictable way. Having a close relative with schizophrenia, bipolar disorder, schizoaffective disorder, or major depression may increase risk. Most relatives of affected people do not develop schizoaffective disorder.

Differences in brain development and chemical signalling may contribute. Research has not identified a single brain scan, chemical imbalance, or gene that explains the condition.

Stressful experiences may affect when symptoms appear or return in someone who is already vulnerable. Stress alone does not fully explain the condition and is not a personal failing.

Cannabis, stimulants like methylphenidate or lisdexamfetamine , hallucinogens, and some other substances can trigger or worsen psychotic or mood symptoms. However, symptoms caused directly by a substance are diagnosed differently. Substance use can also make it harder to understand the pattern of an illness.

Diagnosis

There is no blood test, brain scan, or questionnaire that confirms schizoaffective disorder. Diagnosis is based on a detailed mental health assessment and the pattern of symptoms over time.

A mental health professional will ask about psychotic symptoms, depression, mania, sleep, daily functioning, medicines, alcohol and drug use, physical health, and family history. With permission, information from someone who knows you well may help establish the timeline.

An important feature is a period of hallucinations or delusions lasting at least two weeks without a major depressive or manic episode. Major mood episodes must also be present for most of the overall illness. This pattern may only become clear after repeated assessments.

The diagnosis can change as more information becomes available. This does not necessarily mean that an earlier assessment was careless. Schizophrenia, bipolar disorder with psychotic features, severe depression, trauma-related conditions, and substance-related psychosis can

look similar at first.

A physical examination and tests may be used to look for other causes of symptoms. Depending on the situation, these may include blood tests, urine drug testing, or brain imaging. Conditions affecting the thyroid, nervous system, immune system, or brain can sometimes cause similar changes.

Treatment and management

Treatment is planned around the type and severity of symptoms, previous responses, physical health, side effects, and personal goals. It often combines medicine, psychological therapy, and practical support.

Antipsychotic medicine is usually the main treatment for hallucinations, delusions, and disorganised thinking. It may be taken by mouth or, for some medicines, as a long-acting injection. Regular review is important because side effects can include sleepiness, movement problems, sexual difficulties, weight gain, and changes in blood sugar or cholesterol.

A mood stabilising medicine may be considered when mania is part of the condition. Antidepressants like sertraline or fluoxetine may be used for significant depressive symptoms, usually alongside careful monitoring because they can sometimes contribute to mood elevation in susceptible people.

Do not stop psychiatric medicine suddenly without clinical advice. Sudden changes can lead to withdrawal effects or a return of symptoms. If side effects are difficult, a prescriber can discuss adjustments or alternatives.

Talking therapies can help you understand symptoms, manage stress, improve coping skills, and recognise early warning signs. Cognitive behavioural therapy for psychosis focuses on reducing distress and improving function rather than arguing about whether an experience is real.

Family education or therapy can improve communication and help relatives respond constructively. Rehabilitation services may support education, employment, housing, finances, and daily routines. Treatment for alcohol or drug use can be provided at the same time.

Hospital care may be needed during a severe episode if a person cannot safely meet basic needs, is at immediate risk, or needs close observation while treatment begins. The aim is safety and stabilisation, not punishment.

Electroconvulsive therapy, or ECT, is sometimes considered for severe depression, mania, catatonia, or symptoms that have not improved with other treatment. Catatonia involves major changes in movement and response, such as remaining motionless or barely speaking.

Complications

Untreated or poorly controlled symptoms can disrupt work, study, housing, finances, and relationships. Problems with planning or motivation may also make meals, hygiene, appointments, and medicine routines harder to manage.

Depression and psychosis can increase the risk of self-harm or suicide. Substance use may worsen symptoms, increase impulsive behaviour, or interfere with treatment.

Physical health also matters. The condition itself, reduced activity, smoking, limited access to care, and some medicines can contribute to weight, heart, or metabolic problems. Metabolic problems include high blood sugar and unhealthy cholesterol levels. Routine monitoring can identify these changes early.

During mania or psychosis, judgement may be affected. This can lead to unsafe spending, driving, sexual activity, conflict, or vulnerability to exploitation. Advance planning can help trusted people respond before problems become severe.

Prevention

Schizoaffective disorder cannot currently be reliably prevented. There is no proven way to remove an inherited or biological vulnerability.

It may be possible to reduce relapses and their impact. Helpful measures can include taking agreed treatment, attending follow-up appointments, keeping a regular sleep pattern, avoiding recreational drugs, and seeking help when early symptoms return.

A written relapse plan can record your early warning signs, preferred services, medicine information, and who may be contacted. Common warning signs include sleeping less, becoming unusually suspicious, withdrawing from others, or having rapidly changing energy and mood.

These steps reduce risk but cannot guarantee that another episode will not happen.

Living with it and what to expect

Schizoaffective disorder is usually a long-term condition, but its course is not identical for everyone. Some people have separate episodes with long periods of recovery. Others have ongoing symptoms that rise and fall.

Diagnosis and treatment may take time because the pattern of psychosis and mood

symptoms must be observed across months or years. Finding a medicine that helps without unacceptable side effects may also require several changes.

Recovery does not always mean that every symptom disappears. It may mean returning to study or work, maintaining relationships, living more independently, or learning how to manage voices and mood changes with less distress.

Regular follow-up remains useful when symptoms are stable. Appointments can review sleep, mood, psychosis, physical health, medicine effects, substance use, and personal goals. Blood pressure, weight, blood sugar, cholesterol, and movement symptoms may be monitored when relevant to treatment.

Support from family, friends, peer groups, housing services, or workplace accommodations can make daily life more manageable. A predictable routine and steady sleep often help, especially for people who experience mania.

Setbacks can occur. They do not erase earlier progress. Prompt support when warning signs appear may shorten an episode or reduce its effect.

When to get medical help

Arrange a routine appointment if you notice returning voices, suspicious thoughts, depression, rising energy, reduced sleep, or problems managing treatment. Contact the service sooner if symptoms are changing quickly or interfering with eating, hygiene, work, study, or relationships.

Seek urgent same-day help if you or someone close to you is becoming severely confused, highly agitated, unable to care for basic needs, or rapidly more unwell. Urgent assessment is also appropriate for several nights with almost no sleep alongside increasing energy, impulsiveness, or psychotic symptoms.

Use emergency services if there is:

If it is safe, stay with the person, reduce access to weapons or large quantities of medicine, and share relevant medicine or crisis-plan information with emergency responders.

- An immediate risk of suicide, self-harm, or harm to someone else
- A plan or intention to act on harmful thoughts
- A voice giving harmful commands that feels difficult to resist
- Severe immobility, unresponsiveness, or inability to eat or drink
- Fever, marked muscle stiffness, and confusion while taking antipsychotic medicine

Common questions

Is schizoaffective disorder the same as schizophrenia?

No. Both can involve psychosis, but schizoaffective disorder also includes major mood episodes that are present for most of the illness. The distinction depends on the timing of symptoms and may take time to establish.

Is schizoaffective disorder a type of bipolar disorder?

Bipolar-type schizoaffective disorder includes mania, but it is not the same diagnosis as bipolar disorder. In bipolar disorder with psychotic features, psychosis occurs during a mood episode. In schizoaffective disorder, psychosis also occurs without a major mood episode.

Can schizoaffective disorder go away?

There is no established cure. Some people have long periods with few or no symptoms, while others need continuing treatment and support. Stopping treatment because you feel better can increase relapse risk, so changes should be planned with the prescribing team.

Can people with schizoaffective disorder work or study?

Yes. Many people work, study, volunteer, care for family, or pursue other goals. Flexible hours, gradual returns, reduced workload, quiet spaces, and disability accommodations may help during recovery.

Does schizoaffective disorder make someone violent?

Most people with this condition are not violent. Risk may rise during severe untreated symptoms, particularly when combined with substance use or a history of violence. People with serious mental illness are also vulnerable to being harmed or exploited.

Will I need medicine for life?

Some people benefit from long-term medicine, while treatment plans change for others. The decision is based on previous episodes, relapse risk, benefits, side effects, and preferences. Any reduction should be gradual and clinically supervised.

Sources

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders: DSM-5-TR . Fifth edition, text revision. American Psychiatric Association Publishing; 2022.

- World Health Organization. Clinical Descriptions and Diagnostic Requirements for ICD-11 Mental, Behavioural and Neurodevelopmental Disorders . World Health Organization; 2024.
- American Psychiatric Association. The American Psychiatric Association Practice Guideline for the Treatment of Patients With Schizophrenia . Third edition. American Psychiatric Association Publishing; 2020.
- National Institute for Health and Care Excellence. Psychosis and Schizophrenia in Adults: Prevention and Management . Clinical guideline CG178; 2014.
- MedlinePlus. Schizoaffective Disorder . U.S. National Library of Medicine.
- Malaspina D, Owen MJ, Heckers S, and colleagues. "Schizoaffective Disorder in the DSM-5." Schizophrenia Research . 2013;150(1):21-25.