CONDITION

Simple pulmonary eosinophilia

Simple pulmonary eosinophilia is a temporary lung reaction involving eosinophils, often caused by parasites or medicines.

WHAT IT IS
Temporary eosinophil-related inflammation and patches in the lungs

BODY SYSTEM
Lungs and breathing

MAIN SYMPTOMS
Dry cough, fever, wheezing, shortness of breath

WHAT CAUSES IT
Parasite migration, medicine reactions, or an unknown trigger

HOW IT IS DIAGNOSED
Blood eosinophils, chest imaging, and cause-specific tests

MAIN TREATMENT
Trigger removal and cause-specific treatment

CAN IT BE CURED?
Yes - usually resolves when trigger ends

IS IT AN EMERGENCY?
Sometimes - severe breathing trouble or blue lips

LONG-TERM OUTLOOK
Usually complete recovery without lasting lung damage

Overview

Simple pulmonary eosinophilia is a temporary condition in which eosinophils collect in the lungs. Eosinophils are white blood cells involved in allergic reactions and the body's response to some parasites.

The inflammation can produce hazy or patchy areas on a chest X-ray or CT scan. These patches may move from one part of the lungs to another before disappearing.

The condition is usually mild. Some people have no symptoms and learn about it after a blood test or chest image done for another reason. When symptoms occur, they often resemble a mild chest infection or asthma flare.

Simple pulmonary eosinophilia matters because similar findings can occur in more persistent

conditions. These include chronic eosinophilic pneumonia, certain blood disorders and inflammation of the blood vessels. A clinician will look at how long the changes last, how unwell you are and whether another organ is affected.

Unlike chronic forms of eosinophilic lung inflammation, simple pulmonary eosinophilia usually settles within several weeks and does not normally scar the lungs.

Symptoms

Symptoms are usually mild and may last for only days or weeks. You might have:

Some people cough up a small amount of mucus. Coughing up blood is not typical and needs prompt medical assessment.

Symptoms related to the underlying cause may also occur. A parasite affecting the digestive system can cause stomach pain, nausea or changes in bowel habits. A medicine reaction may cause a rash, itching or fever.

The number of eosinophils in the blood does not always match the severity of symptoms. A high count can occur with few breathing problems, while significant breathlessness needs assessment regardless of the blood result.

- A dry cough
- A mild fever
- Wheezing or chest tightness
- Shortness of breath
- Tiredness or a general unwell feeling

Causes and risk factors

The condition is often triggered by parasite larvae passing through the lungs during part of their life cycle. Parasites associated with this pattern include roundworm, hookworm and Strongyloides . Exposure is more likely in places where these parasites are common and sanitation is limited.

Risk can be affected by:

Some medicines can trigger eosinophilic inflammation in the lungs. Several types have been linked to this reaction, including certain antibiotics and anti-inflammatory medicines. The reaction may begin soon after starting a medicine, but timing is not always predictable.

Do not stop a prescribed medicine on your own unless you are having signs of a severe allergic reaction. Contact the prescriber promptly so the possible trigger and alternatives can be reviewed.

Sometimes no cause is found. This is called idiopathic, meaning that the trigger remains unknown after an appropriate evaluation.

Having asthma or allergies can raise the eosinophil count for other reasons. These conditions do not by themselves prove that you have simple pulmonary eosinophilia.

- Living in or travelling to an area where these parasites occur
- Contact with soil contaminated by human waste
- Walking barefoot on contaminated soil
- Eating food contaminated with parasite eggs

Diagnosis

There is no single test that confirms every case. Diagnosis is based on the combination of symptoms, blood tests, chest imaging and the way the condition changes over time.

A complete blood count may show an increased number of eosinophils. This is called peripheral eosinophilia. A normal count does not completely rule out eosinophilic lung inflammation, particularly early in the illness.

A chest X-ray may show temporary, patchy areas of inflammation. A CT scan gives more detailed images and may be used when the X-ray is unclear, symptoms are more significant or another condition needs to be excluded.

Your evaluation may include questions about:

Stool tests or blood tests may be used when a parasite is possible. More than one stool sample can sometimes be needed because parasites are not released continuously. The most useful test depends on the suspected parasite and the stage of infection.

Tests of mucus, lung function or, less commonly, a bronchoscopy may be needed if the diagnosis remains uncertain. Bronchoscopy uses a thin flexible camera to examine the airways and collect samples.

Simple pulmonary eosinophilia becomes less likely when symptoms are severe, lung changes persist, or the heart, nerves, skin, kidneys or other organs are affected. In that situation, clinicians investigate other causes of pulmonary eosinophilia.

- Recent and past travel or residence

- Contact with soil or sanitation risks
- New medicines, supplements and non-prescription products
- Asthma, allergies and previous eosinophil results
- Rashes, numbness, digestive symptoms or other organ problems

Treatment and management

Many mild cases improve without treatment once the trigger has passed. Follow-up is still important to make sure symptoms, eosinophil levels and chest changes are resolving.

If a parasite is identified or strongly suspected, treatment targets that specific infection. The choice of antiparasitic medicine depends on the organism, test results and your health circumstances.

If a medicine is the likely cause, a clinician may advise stopping it or changing to an alternative. The suspected reaction should be recorded in your medical notes. Not every symptom after starting a medicine is an allergy, so the timing and other possible causes need careful review.

Corticosteroids like prednisone or hydrocortisone reduce inflammation but are not routinely needed for uncomplicated, improving simple pulmonary eosinophilia. Persistent or serious symptoms may suggest a different condition and need specialist assessment.

Possible Strongyloides infection is especially important before corticosteroids or other immune-suppressing treatment. These medicines can allow that parasite to multiply rapidly and cause a serious widespread infection. Tell your healthcare team about relevant travel or residence, even if it occurred many years ago.

Treatment may also include supportive care for symptoms, such as managing wheezing. This should be based on your examination and medical history rather than used as proof of the diagnosis.

Complications

Complications are uncommon when the condition is truly simple pulmonary eosinophilia. Lasting lung damage and scarring are not typical.

Problems can occur if the underlying trigger remains. A parasite may continue to cause symptoms elsewhere in the body, while continued exposure to a responsible medicine can lead to renewed inflammation.

The main concern is that persistent or worsening findings may represent another condition. Chronic eosinophilic pneumonia, allergic reactions to fungi, inflammation of blood vessels and disorders that produce very high eosinophil counts can require different treatment.

A parasite infection can also become more serious in someone whose immune system is weakened. Strongyloides is a particular concern because it can remain in the body for years and become dangerous during immune-suppressing treatment.

Follow-up testing helps confirm that the expected recovery is happening rather than allowing a more persistent cause to go unnoticed.

Prevention

Simple pulmonary eosinophilia cannot always be prevented. Medicine reactions may be unpredictable, and no specific trigger is found in some cases.

Measures that can reduce exposure to certain parasites include wearing shoes where soil contamination is possible, using safe water, washing produce and following local food-safety advice. These steps reduce risk but cannot remove it completely.

If you have previously had a suspected medicine-related episode, keep a record of the medicine and the reaction. Share it when receiving new prescriptions. Do not label unrelated medicines as unsafe without a medical review, because this can unnecessarily limit future treatment choices.

Tell clinicians about previous residence or long-term travel in places where Strongyloides occurs, especially before treatment that suppresses the immune system. An infection can remain silent for a long time.

Living with it and what to expect

Symptoms often settle over days to several weeks. Chest X-ray changes generally clear within about a month, although the exact timing is not the same for everyone.

Your clinician may repeat a blood count or chest image to confirm improvement. Routine follow-up is particularly useful when the original eosinophil count was high, a parasite or medicine was suspected, or the first image showed clear lung changes.

Most people do not need long-term breathing treatment once the episode has resolved. Simple pulmonary eosinophilia does not usually limit exercise or everyday activities in the long term.

During recovery, return to activity according to how you feel. Breathlessness that is getting worse rather than better should not be treated as a normal part of recovery.

If symptoms or imaging changes continue beyond the expected short course, the diagnosis may need to be reconsidered. This does not automatically mean there is a serious problem, but it may lead to further blood tests, repeat imaging or referral to a lung specialist.

Recurrence is possible if the original trigger is encountered again or if an underlying parasite has not been fully addressed. When no cause is found and all findings resolve, repeated episodes are not expected in most people.

When to get medical help

Arrange a routine medical appointment if you have an unexplained cough, wheezing or fever that is not improving, particularly if you have recently started a medicine or have relevant travel or soil exposure.

Seek prompt medical advice if:

Call emergency services for severe difficulty breathing, blue or grey lips, fainting, confusion, or chest pain with marked breathlessness. These symptoms have many possible causes and should be assessed urgently rather than assumed to be simple pulmonary eosinophilia.

- Breathing symptoms are worsening
- Fever is persistent or you feel increasingly unwell
- You cough up blood
- Symptoms return after previously improving
- You develop a widespread rash, swelling or new numbness
- Follow-up tests remain abnormal

Common questions

Is simple pulmonary eosinophilia cancer?

No. It is an inflammatory reaction involving eosinophils, not a cancer. However, unusual or persistent eosinophil findings sometimes require tests to exclude blood or immune-system disorders.

Is it contagious?

The lung inflammation itself is not contagious. Some parasite infections can spread through

contaminated soil, food or water, but they are not usually passed to another person by coughing.

Will it go away on its own?

Many cases resolve without anti-inflammatory treatment after the trigger has passed. An identified parasite or ongoing medicine reaction may still need specific management. Follow-up confirms that the lungs and blood results are returning to normal.

How long does recovery take?

Symptoms often improve within days or weeks, and lung patches commonly disappear within about a month. Continued symptoms or abnormal images need reassessment because they are less typical of the simple form.

Can it come back?

It can return if you are exposed to the same trigger again or if an underlying infection remains. If no trigger is found and all findings clear, long-term recurrence is not typical.

Is it the same as pneumonia?

It can look like pneumonia on a chest image, but it is not usually caused by ordinary pneumonia bacteria. The patches are produced by eosinophilic inflammation and may move or disappear relatively quickly.

Sources

- Merck Manual Professional Edition: Simple Pulmonary Eosinophilia
- Rosenberg CE, Khoury P. Approach to Eosinophilia Presenting With Pulmonary Symptoms. Chest . 2021;159(2):507-516
- Cottin V. Eosinophilic Lung Diseases. Clinics in Chest Medicine . 2016;37(3):535-556
- Centers for Disease Control and Prevention: DPDx-Ascariasis
- Centers for Disease Control and Prevention: DPDx-Hookworm
- Centers for Disease Control and Prevention: Clinical Overview of Strongyloides