CONDITION

Skeletal limb abnormalities

Skeletal limb abnormalities are differences in the bones or joints of an arm or leg.

WHAT IT IS
Structural differences in arm or leg bones and joints

BODY SYSTEM
Bones, joints, muscles, and limbs

MAIN SYMPTOMS
Altered limb shape, length difference, limited movement, uneven walking

WHAT CAUSES IT
Genetic changes, development differences, injury, or bone conditions

HOW IT IS DIAGNOSED
Examination and X-rays; sometimes prenatal imaging or genetic testing

MAIN TREATMENT
Monitoring, therapy, braces, prostheses, or surgery

CAN IT BE CURED?
Sometimes - correction is possible for selected differences

IS IT AN EMERGENCY?
Sometimes - sudden severe pain or loss of circulation

LONG-TERM OUTLOOK
Many people live active lives; support needs differ

Overview

Skeletal limb abnormalities are differences in the bones or joints of an arm or leg. They may affect the limb's length, shape, alignment, movement, or number of bones. Some are present at birth. Others develop after an injury, infection, growth problem, or another condition.

This is a broad category, not one diagnosis. Examples include a shortened or missing part of a limb, extra or joined fingers or toes, bowed bones, unequal leg lengths, and joints that did not form in the usual way. A difference may affect one limb or several limbs.

Some skeletal limb abnormalities mainly change appearance. Others affect walking, balance, grip, reach, or everyday tasks. Pain is not always present. A person with a substantial visible difference may have little pain, while a smaller alignment problem can sometimes cause discomfort or difficulty with movement.

These differences also range from isolated findings to parts of genetic syndromes that affect other areas of the body. Identifying the specific abnormality helps clarify whether other checks, treatment, or long-term monitoring may be useful.

Symptoms

The signs depend on which bones and joints are affected. They may be noticed during pregnancy, at birth, when a child begins using the limb, or later during growth.

Possible signs include:

A mild leg-length difference may cause no clear symptoms. A larger difference can lead to limping, toe walking on one side, or a tilted pelvis. During growth, the difference may become more noticeable even when the underlying condition is unchanged.

New pain, swelling, numbness, or loss of function should not automatically be assumed to come from a known limb difference. These symptoms may have another cause and deserve assessment.

- A limb, hand, foot, finger, or toe with an unusual shape
- One arm or leg being shorter than the other
- A missing, shortened, extra, curved, or joined bone or digit
- A joint that moves less than expected or feels unstable
- Uneven shoulders, hips, standing posture, or walking
- Difficulty gripping, reaching, running, climbing stairs, or using footwear
- Tiredness or pain after activity
- Repeated pressure areas, blisters, or skin irritation from braces or prostheses

Causes and risk factors

Congenital limb abnormalities develop before birth. "Congenital" means present at birth, although the difference may not be recognised until later.

Some congenital differences result from a change in a gene or chromosome. They may run in families, but many occur in a child with no family history. A genetic change can affect only the limbs or be part of a syndrome involving other organs.

Other differences happen when a limb does not develop in the usual sequence during

pregnancy. Rarely, strands from the amniotic sac surrounding the fetus can restrict a developing limb. Certain infections, medicines, or harmful exposures during pregnancy are linked with particular abnormalities, but the cause often remains unknown. In most cases, the difference is not caused by something a parent did.

Skeletal limb abnormalities can also develop after birth. Possible causes include:

A family history or a known genetic condition may increase the chance of some congenital differences. The chance of an acquired abnormality is linked more closely to the underlying injury or condition.

- Injury to a bone's growth plate, where childhood bone growth occurs
- Bone or joint infection
- Conditions affecting normal bone growth
- Reduced blood supply to part of a bone
- Neuromuscular conditions that alter muscle balance and joint position
- A bone tumour or its treatment
- Major trauma or surgery involving a limb

Diagnosis

Assessment usually begins with a medical history and physical examination. The clinician may ask when the difference was first noticed, whether it has changed, and how it affects movement or daily activities.

The examination may compare the length, alignment, strength, sensation, and joint movement of both sides. For a leg difference, walking and posture may be assessed. Measurements may be repeated as a child grows because future growth can affect treatment decisions.

X-rays show the number, shape, alignment, and maturity of bones. Standing X-rays or specialised length measurements may be used for unequal legs. Ultrasound, CT, or MRI may be needed when more detail about cartilage, joints, muscles, nerves, or blood vessels is important. CT uses X-rays to create detailed images, while MRI uses a magnetic field.

Some congenital abnormalities can be seen on a pregnancy ultrasound. Prenatal imaging cannot always show how well a limb will function after birth, so further assessment is usually needed later.

Genetic testing may be offered when the pattern suggests an inherited condition or syndrome. This may involve a blood or saliva sample. A genetics appointment can explain what the result

may mean for the person and their family. Not every limb difference has an identifiable genetic cause.

Treatment and management

Treatment is based on function, symptoms, age, growth remaining, and the type of bone or joint difference. Appearance may also matter to the person, but treatment does not need to make a limb look typical to be successful.

A mild difference that does not cause pain or limit activity may only need observation. Children may have regular measurements or X-rays to check how the limb changes with growth.

Physical therapy can support strength, movement, balance, and walking. Occupational therapy focuses on daily activities such as dressing, writing, preparing food, or using equipment. Therapy may also help someone find ways of completing tasks that suit their body.

Braces, splints, shoe lifts, or specially made footwear may improve alignment, stability, or comfort. A prosthesis is an artificial limb or limb part. Some people use one for selected activities, while others prefer not to use one. Prosthetic needs can change with growth, activity, comfort, and personal preference.

Surgery may be considered when it is likely to improve function, alignment, comfort, or limb stability. Depending on the abnormality, surgery may:

Limb lengthening is a lengthy process that gradually separates a surgically divided bone so new bone can form in the gap. It requires close follow-up and rehabilitation and carries risks such as infection, stiffness, nerve problems, and delayed bone healing.

Pain management may include activity changes, therapy, supportive equipment, or medicines. The appropriate approach depends on the reason for the pain rather than the limb's appearance alone.

Emotional support can also be important. Children and adults may face unwanted attention, teasing, body-image concerns, or frustration with inaccessible environments. Peer groups and limb-difference organisations can provide practical and social support.

- Release joined fingers or toes
- Reconstruct or stabilise a joint
- Straighten a curved bone
- Slow growth in the longer leg
- Shorten a bone or gradually lengthen a limb

- Remove an extra digit or a painful non-functional structure
- Prepare a limb for more effective prosthetic use

Complications

Possible complications depend on the specific abnormality and its treatment. They can include reduced joint movement, muscle weakness, unstable joints, uneven walking, or pain from altered pressure on bones and joints.

Over time, unusual joint loading may contribute to earlier wear of a joint, called osteoarthritis. A leg-length difference can affect the hips, knees, back, or energy used while walking, although small differences do not always cause problems.

Braces and prostheses can cause pressure areas if the fit changes. Growing children need regular fit checks. Surgery can lead to infection, stiffness, scarring, problems with bone healing, or injury to nearby nerves and blood vessels.

Not everyone develops complications. Monitoring is intended to identify meaningful changes without treating differences that are not causing a problem.

Prevention

Many congenital skeletal limb abnormalities cannot be prevented. This is especially true when the cause is an inherited or newly occurring genetic change. Even when no cause is found, it does not mean that a parent caused the difference.

General pre-pregnancy and prenatal care may reduce some health risks but cannot guarantee typical limb development. This includes reviewing medicines before pregnancy, avoiding tobacco and harmful substances, and managing existing health conditions. Prescribed medicine should not be stopped without medical advice.

Some acquired abnormalities may be less likely with injury prevention and prompt care for serious bone, joint, or growth-plate injuries. However, infections, growth disorders, and other causes are not always preventable.

Living with it and what to expect

The long-term course is different for each type of limb abnormality. A congenital difference does not usually spread to other limbs. However, its effect can change as the body grows,

activities become more demanding, or joints experience years of altered movement.

Some children need little more than periodic review. Others have therapy, equipment changes, or several operations spaced across childhood and adolescence. Growth-related treatment usually becomes less frequent after the skeleton matures, although adults may still need care for pain, joint wear, prosthetic fit, or previous surgery.

Many people with limb differences attend mainstream school, work, drive, exercise, play sports, and live independently. Adapted tools, changes to the home or workplace, and suitable sports equipment can make activities easier. The most useful adaptation is the one that supports the person's own goals.

Routine follow-up may include checking growth, joint movement, skin health, walking, equipment fit, and the effect on daily life. A new stage of life, such as starting school, changing jobs, pregnancy, or taking up a sport, may create different needs without meaning that the abnormality has worsened.

When to get medical help

Arrange a routine appointment if you notice a persistent difference in limb length, shape, joint movement, or walking. Assessment is also appropriate if an existing difference begins causing pain, limits activity, or affects footwear, braces, or a prosthesis.

Seek prompt medical advice for:

Get emergency help after an injury or procedure if the limb becomes pale, blue, unusually cold, or difficult to move. Rapidly worsening severe pain, major swelling, loss of sensation, uncontrolled bleeding, or an open injury with visible bone also requires emergency assessment. These signs can indicate reduced blood flow, nerve injury, compartment syndrome, or a serious fracture.

- New redness, warmth, swelling, drainage, or fever
- Increasing pain or reduced movement without a clear reason
- A brace, splint, or prosthesis causing sores or broken skin
- A cast that feels too tight, becomes damaged, or gets wet inside
- A child suddenly refusing to use a limb or bear weight
- New numbness, weakness, or repeated falls

Common questions

Is a skeletal limb abnormality a disability?

Not always. Disability is based on how a condition affects daily activities, not only on appearance or a diagnosis. Some people need no assistance. Others may qualify for accommodations, equipment, therapy, or disability support because movement or hand function is substantially affected.

Can a limb abnormality be seen on a pregnancy ultrasound?

Some can be seen, especially major differences in limb length, position, or formation. Smaller bone or joint differences may not be visible. Ultrasound findings may lead to more detailed prenatal imaging or genetic assessment, but function often cannot be fully predicted before birth.

Will my child need surgery?

Not necessarily. Surgery is usually considered when it may improve function, alignment, stability, comfort, or future growth. Some differences are best managed with observation, therapy, or equipment. Decisions may be revisited as a child grows.

Do limb-length differences get worse with age?

A difference caused by altered growth can increase while a child is growing. Clinicians may estimate future length by looking at current measurements, bone maturity, and the underlying cause. Once growth has finished, the bone-length difference itself is usually stable, although its effect on joints or walking may change.

Can someone with a limb difference play sports?

Often, yes. The most suitable activities and adaptations depend on strength, joint stability, skin health, balance, and personal interest. Some people use a sport-specific prosthesis or brace. Pain, repeated skin injury, or joint instability should be assessed rather than pushed through.

Is it genetic, and can it happen again in a family?

Some limb abnormalities are genetic, but many are not inherited and have no identified cause. The chance of recurrence depends on the exact diagnosis and any genetic test result. Genetic counselling can provide a family-specific explanation without assuming that every limb difference is inherited.

Sources

- Centers for Disease Control and Prevention. About Limb Reduction Defects , updated May 15, 2024
- World Health Organization. Congenital disorders , fact sheet, February 27, 2023
- American Academy of Orthopaedic Surgeons OrthoInfo. Lower Limb Length Discrepancy
- MedlinePlus Medical Encyclopedia. Congenital limb defects , U.S. National Library of Medicine
- Farr S, Catena N, Martínez-Álvarez S, Soldado F. Peromelia-congenital transverse deficiency of the upper limb: a literature review and current prosthetic treatment . Journal of Children's Orthopaedics . 2018;12(6):558-565
- Sabharwal S, Kumar A. Methods for assessing leg length discrepancy . Clinical Orthopaedics and Related Research . 2008;466:2910-2922