CONDITION

Stoddard solvent poisoning

Stoddard solvent poisoning is illness caused by breathing, swallowing, or having significant contact with this petroleum-based solvent.

WHAT IT IS
Harmful exposure to a petroleum-based cleaning and thinning solvent

BODY SYSTEM
Lungs, nervous system, skin, and eyes

MAIN SYMPTOMS
Cough, breathing difficulty, dizziness, nausea

WHAT CAUSES IT
Breathing, swallowing, or skin contact with petroleum solvent

HOW IT IS DIAGNOSED
Exposure history, examination, oxygen measurement, and chest X-ray

MAIN TREATMENT
Decontamination, breathing support, monitoring, and other supportive care

CAN IT BE CURED?
Yes - most acute cases resolve with supportive care

IS IT AN EMERGENCY?
Sometimes - breathing trouble, collapse, confusion, or seizures

LONG-TERM OUTLOOK
Most recover; severe lung injury can have lasting effects

Overview

Stoddard solvent is a refined petroleum product used for cleaning, degreasing, dry cleaning, and thinning some paints and coatings. It is also called a type of mineral spirit or white spirit, although products sold under those names may contain different chemical mixtures.

Poisoning can happen after breathing concentrated fumes, swallowing the liquid, or having heavy or repeated contact with the skin. The most important risk after swallowing is aspiration . This means the liquid enters the lungs, often during swallowing or vomiting. Even a small amount in the lungs can cause chemical inflammation called hydrocarbon pneumonitis.

Breathing high concentrations can affect the brain and nervous system. It may cause headache, dizziness, poor coordination, confusion, or loss of consciousness. The solvent can also irritate the eyes, airways, and skin.

A brief, low-level exposure may cause only temporary irritation. More serious exposure can interfere with breathing and may require hospital care. If an exposure has just happened, a poison information service can give advice based on the product, route of exposure, symptoms, and amount involved.

Symptoms

Symptoms depend on how the exposure happened and how concentrated it was.

Breathing Stoddard solvent fumes may cause:

After swallowing, there may be coughing, choking, gagging, nausea, vomiting, or stomach discomfort. Coughing soon after swallowing can be a sign that some liquid entered the airways.

Lung symptoms may not be obvious immediately. Cough, rapid breathing, chest pain, fever, low blood oxygen, or increasing breathlessness can develop over the following hours if aspiration has occurred. Fever after aspiration can result from chemical inflammation and does not always mean there is an infection.

Eye contact can cause pain, redness, watering, or blurred vision. Skin contact often causes dryness, redness, cracking, or a burning feeling. Repeated contact removes the skin's protective oils and may lead to dermatitis, which is inflamed, itchy skin.

Very high exposure can cause seizures, an abnormal heart rhythm, coma, or stopped breathing. These effects are uncommon but require emergency care.

- Burning or irritation in the nose and throat
- Headache, dizziness, nausea, or unusual tiredness
- Cough, chest tightness, or shortness of breath
- Poor balance, confusion, drowsiness, or fainting after heavy exposure

Causes and risk factors

Stoddard solvent poisoning is caused by enough of the chemical entering the body to cause harm. The main routes are inhalation, swallowing, and skin or eye contact.

Inhalation exposure is more likely when the solvent is used in a small or poorly ventilated area. Spraying, heating, or using large open containers may increase the amount in the air. Entering a heavily contaminated space without suitable protective equipment can be especially dangerous.

Swallowing may happen when the solvent is stored in a cup, drink bottle, or unlabelled container. Young children may swallow it accidentally. Intentional inhalation or swallowing can lead to severe poisoning.

People who work in painting, machinery maintenance, printing, metal cleaning, dry cleaning, or similar jobs may have repeated exposure. Risk increases when ventilation and protective measures are inadequate.

Damaged skin may absorb more solvent than intact skin, but widespread poisoning through the skin alone is less common. Long periods in solvent-soaked clothing can still cause significant irritation and exposure.

Alcohol, sedatives, or other substances that reduce alertness may add to the solvent's effects on the brain. They may also make choking or aspiration more likely.

Diagnosis

There is no single blood test that confirms Stoddard solvent poisoning or reliably measures how severe it is. Diagnosis is based mainly on what happened, the product involved, symptoms, and examination findings.

Bring the product container, label, or safety data sheet if this can be done safely. Different products described as mineral spirits or paint thinner may contain different chemicals, so the exact product matters.

A healthcare team may check:

A chest X-ray can be normal soon after exposure even when lung irritation is developing. For this reason, symptoms and oxygen levels are important, and observation may be needed after a concerning ingestion or inhalation.

- Breathing rate, heart rate, blood pressure, temperature, and alertness
- Blood oxygen using a sensor placed on a finger
- Lung sounds and signs of skin or eye injury
- A chest X-ray if aspiration or lung injury is suspected
- An electrocardiogram, or ECG, to check the heart rhythm
- Blood tests when needed to assess oxygen, acid balance, or organ function

Treatment and management

Treatment begins by stopping the exposure without putting anyone else at risk. Do not enter an enclosed, contaminated space unless trained and properly equipped.

After inhalation, move to fresh air if it is safe to do so. Loosen tight clothing and avoid physical exertion. Anyone with breathing difficulty, marked drowsiness, confusion, or collapse needs emergency help.

For skin exposure, remove contaminated clothing and wash the skin thoroughly with soap and water. Contaminated clothing should be handled carefully because it may continue releasing fumes and may be flammable.

For eye exposure, remove contact lenses if they come out easily and rinse the eye continuously with clean, lukewarm water. Persistent pain, redness, light sensitivity, or blurred vision needs medical assessment.

If Stoddard solvent has been swallowed, do not make the person vomit. Vomiting increases the chance of solvent entering the lungs. Do not give food, drink, activated charcoal, or another home remedy unless a poison specialist or healthcare professional advises it. Contact a poison information service promptly.

Hospital treatment is supportive, meaning it protects breathing and other body functions while the chemical's effects wear off. It may include:

Activated charcoal is generally not useful for Stoddard solvent and may increase aspiration risk. Procedures intended to empty the stomach are also generally avoided because of that risk.

Antibiotics like amoxicillin or doxycycline do not treat chemical lung inflammation itself. They may be used if there is evidence of a separate bacterial infection. Steroids like prednisone or hydrocortisone are not routinely used for uncomplicated hydrocarbon pneumonitis.

- Supplemental oxygen and repeated oxygen measurements
- Medicines to open narrowed airways when wheezing is present
- Fluids and treatment for low blood pressure when needed
- Breathing support, including mechanical ventilation in severe cases
- Heart monitoring after significant exposure
- Treatment of seizures or other complications

Complications

The main complication is hydrocarbon pneumonitis, a chemical injury to the lungs caused by

aspiration. It can lead to low blood oxygen, fluid and inflammation in the lungs, or respiratory failure.

Other possible complications include:

Repeated high workplace exposure has also been linked with ongoing headaches, problems with concentration, mood changes, and reduced coordination. These symptoms can have other causes, so they need a careful occupational and medical assessment.

- Abnormal heart rhythms after heavy exposure
- Seizures, coma, or brain injury if oxygen levels become very low
- Persistent dermatitis after repeated skin contact
- Eye surface injury from prolonged contact
- Lasting breathing problems after severe lung damage

Prevention

Most accidental Stoddard solvent poisoning can be prevented by controlling exposure.

Use the product only as directed and with effective ventilation. Indoor use may require local exhaust equipment rather than an open window alone. Keep containers closed when they are not being used.

Store the solvent in its original, clearly labelled container. Never transfer it into a water bottle, cup, food container, or other packaging that could be mistaken for a drink. Keep it away from children and from heat, sparks, and flames.

At work, follow the safety data sheet and workplace chemical-control plan. Gloves, eye protection, protective clothing, or respiratory protection may be needed for some tasks. The correct equipment depends on the product and exposure level. Respirators should be selected, fitted, and maintained through a proper workplace respiratory-protection program.

Wash exposed skin rather than cleaning it with more solvent. Change out of contaminated clothing promptly. Do not eat, drink, smoke, or vape in areas where the solvent is handled.

Substituting a less hazardous product or using a closed cleaning system may reduce repeated occupational exposure.

Living with it and what to expect

Many people with mild irritation or brief inhalation exposure improve after they leave the

contaminated area. Headache, nausea, or dizziness often settles as exposure stops, but symptoms that continue or return should be assessed.

If solvent enters the lungs, coughing and breathing symptoms may worsen during the hours after exposure. Recovery from chemical pneumonitis may take days or weeks. Severe cases can require intensive care, and a small number of people are left with reduced lung function or other lasting effects.

Dry or inflamed skin usually improves when contact stops and the skin barrier recovers. Dermatitis may return if exposure continues, so workplace changes and suitable skin protection can be important.

After substantial repeated exposure, problems with memory, concentration, balance, or mood may improve gradually once exposure ends. Some people have persistent symptoms. Follow-up may involve primary care, occupational medicine, lung testing, or neurological assessment based on the effects experienced.

Routine follow-up is reasonable after significant poisoning, especially if there was lung injury, admission to hospital, or ongoing workplace exposure. New or worsening breathing symptoms are not part of routine follow-up and need prompt assessment.

When to get medical help

Call emergency services now if an exposed person has:

Do not enter a contaminated enclosed area to attempt a rescue unless you have appropriate training and protective equipment.

Contact a poison information service promptly after swallowing Stoddard solvent, even if there are no symptoms yet. In the United States, Poison Control can be reached at 1-800-222-1222. Elsewhere, use your local poison centre or emergency advice service.

Seek urgent medical assessment for increasing cough, fever, chest pain, rapid breathing, wheezing, vomiting, or unusual drowsiness after exposure.

Arrange routine medical follow-up for skin irritation that does not settle, recurring symptoms at work, or ongoing headaches, concentration problems, cough, or breathlessness. Workplace exposures can also be reviewed by an occupational health service.

- Trouble breathing, blue or grey lips, or severe chest tightness
- Collapse, inability to wake, severe confusion, or a seizure
- Choking or repeated coughing after swallowing the solvent

- Severe weakness, poor coordination, or an abnormal heartbeat
- Exposure inside an enclosed space followed by significant symptoms

Common questions

What should I do if I swallowed mineral spirits?

Do not make yourself vomit. Rinse your mouth and contact a poison information service immediately for product-specific advice. Call emergency services if there is coughing, choking, drowsiness, confusion, or breathing difficulty.

Can Stoddard solvent fumes make me sick?

Yes. Concentrated fumes can irritate the airways and affect the brain, causing headache, nausea, dizziness, drowsiness, or poor coordination. Move to fresh air if it is safe. Significant or continuing symptoms need medical advice.

Is Stoddard solvent the same as paint thinner?

Stoddard solvent is a type of petroleum-based mineral spirit and may be used in paint thinners. However, "paint thinner," "mineral spirits," and "white spirit" are broad product terms. Formulations can differ, so check the label or safety data sheet.

Can a small swallowed amount damage the lungs?

It can if the liquid enters the airway. The lung risk is related not only to the amount swallowed but also to coughing, choking, or vomiting. A poison centre should assess any ingestion rather than waiting for symptoms.

Can Stoddard solvent cause permanent damage?

Most people recover from limited acute exposure. Permanent damage is more likely after severe aspiration, prolonged low oxygen, or repeated high exposure. Ongoing breathing, balance, memory, or concentration problems need follow-up.

Should I get a chest X-ray after exposure?

Not every exposure requires one. A chest X-ray may be used when there is coughing, choking, low oxygen, abnormal lung sounds, or concern that solvent entered the lungs. Early X-rays can be normal, so the decision also depends on symptoms and observation.

Sources

- Agency for Toxic Substances and Disease Registry. Toxicological Profile for Stoddard Solvent . US Department of Health and Human Services; 1995.
- National Institute for Occupational Safety and Health. NIOSH Pocket Guide to Chemical Hazards: Stoddard Solvent . Centers for Disease Control and Prevention.
- Manoguerra AS, Cobaugh DJ, and the American Association of Poison Control Centers. "Hydrocarbon Poisoning: An Evidence-Based Consensus Guideline for Out-of-Hospital Management." Clinical Toxicology . 2007;45(6):662-676.
- Merck Manual Professional Edition. "Hydrocarbon Poisoning." Section on injuries and poisoning.
- MedlinePlus Medical Encyclopedia. "Stoddard Solvent Poisoning." US National Library of Medicine.