SLEEP & STRESS

How to Wind Down When Your Mind Will Not

A racing mind needs less pressure, not another perfect bedtime ritual-and persistent sleeplessness deserves more than self-help.

Your body is in bed. Your day is not

The room is quiet, but your mind has reopened the office. An awkward conversation gets a rewrite. Tomorrow's list grows new branches. Then comes the thought that turns wakefulness into a verdict: you should be asleep by now.

This is where winding down can become another job you believe you are failing. You arrange the room, try to relax and wait for sleep to arrive on command. Each wakeful minute seems to raise the stakes. The effort meant to calm you becomes part of the noise.

Stress and poor sleep do travel together, although that does not prove one always causes the other. A 2022 review of 34 studies in undergraduate students found moderate associations between stress and both poor sleep quality and insomnia. The results varied considerably between studies, so they should be read cautiously-and they may not apply neatly beyond students. Still, the pattern will feel familiar if your worries become loudest after the lights go out.

One difficult night is not automatically a disorder. Short-term insomnia can follow work stress, family pressure or a traumatic event and may last days or weeks. But the distinction may offer little comfort at midnight. What helps first is dropping the demand to solve sleep immediately and looking at what your evening is asking your brain to absorb.

Make the landing less abrupt

A wind-down is not a trick that guarantees unconsciousness. It is a softer boundary between the active part of your day and the bed. The National Heart, Lung, and Blood Institute suggests a routine built around quiet activities such as reading, soothing music or a hot bath. Meditation, yoga and massage are other relaxation approaches a healthcare provider may recommend.

The room can do some of the work. NHLBI advises keeping the bedroom cool, quiet and dark. Electronic devices can interfere with the sleep-wake cycle, so putting them away before bed is part of both NHLBI and CDC guidance. That does not mean one late message has ruined your night. It means the final stretch of the evening need not keep delivering fresh material for your

mind to process.

Consistency matters too. Public-health guidance recommends going to bed and getting up at around the same times, including on weekends. Irregular schedules, night work and frequent shifts can contribute to sleep problems, although NHLBI acknowledges that steadiness may be difficult for shift workers and new parents. Advice that ignores your real life is not much of a wind-down.

Most adults need roughly seven to nine hours of sleep, but duration is only part of the picture. Sleep quality matters because repeated interruptions can reduce time in certain sleep stages. Quality sleep is uninterrupted and refreshing. Your goal, then, is not simply to accumulate time in bed while arguing with yourself about whether it counts.

Persistent trouble sleeping is not a failure to wind down correctly.

Notice what keeps reopening the door

Caffeine deserves an honest look, especially when your mind feels alert long after you want the day to end. A 2018 systematic review found that caffeine typically lengthened the time it took to fall asleep, reduced total sleep time and sleep efficiency, and worsened perceived sleep quality. Timing and amount mattered, but people differed markedly in their sensitivity. Most included research involved men in Western countries, limiting how broadly the findings can be applied.

Alcohol can be deceptive in a different way. NHLBI notes that it may make falling asleep easier while producing lighter sleep and making nighttime waking more likely. Nicotine and late caffeine can also interfere. You do not need to turn the evening into a forensic investigation, but it is worth noticing whether the things used to get through the day are following you into the night.

Then there is the nap taken because last night went badly. NHLBI says avoiding naps, particularly later ones, may help you sleep longer at night. Regular daytime physical activity and regular meals are also included in its healthy-sleep guidance, while exercise close to bed can make falling asleep harder. These are possibilities to observe, not a moral scorecard.

A simple sleep diary can make those patterns easier to see. The CDC suggests recording when you go to bed, wake during the night and rise in the morning, along with naps, exercise, alcohol, caffeine and medicines. The diary is not there to grade your sleep. It can give you-and, if needed, your healthcare provider-a clearer account than the exhausted impression that every night is terrible.

Do not let the cure become another worry

When you are tired, a supplement can look like a clean answer. Magnesium is often promoted for sleep, but the evidence is thin. A 2021 review found only three randomized trials involving 151 older adults with insomnia. Participants taking magnesium fell asleep about 17 minutes sooner than those taking placebo, while the increase in total sleep time was not statistically significant. The evidence was rated low to very low quality, and every trial had a moderate-to-high risk of bias.

Melatonin is not a consequence-free shortcut either. NHLBI says supplements can vary in dose and purity because they are not regulated like medicines. Reported side effects include excess sleepiness, headaches, stomach upset, blood-pressure changes and worsening depression symptoms. Supplements can also interact with medicines. Talk with a healthcare provider before using one, particularly if you are pregnant or trying to become pregnant.

Prescription medicines can help some adults with insomnia, but they require a risk-benefit conversation rather than guesswork. A 2023 network meta-analysis of 69 studies involving 17,319 patients found meaningful differences in effectiveness and tolerability among drug classes. Some performed better than placebo on measures such as falling asleep or time awake after sleep began. Yet the long-term adverse effects of all the medications remained unclear.

This is why a racing mind should not send you straight from frustration to a shopping basket. Most sleep medicines are generally intended for short-term use, and many have side effects. If habits and relaxation are not enough, the next step may be a fuller conversation about the problem-not a stronger promise on a bottle.

Know when the night is asking for help

Persistent trouble sleeping is not a failure to wind down correctly. Chronic insomnia lasts a month or longer, and it may be related to another medical condition, a medicine, another sleep disorder, depression or anxiety. Sometimes the cause is not understood. Cognitive behavioral therapy for insomnia is generally recommended as the first option for long-term insomnia, according to NHLBI, and it can be delivered in person, by telephone or online.

CBT-I is more than generic relaxation. It addresses nervousness about not sleeping, teaches sleep habits and helps rebuild the link between bed and sleep. One element involves leaving bed when you cannot sleep rather than letting it become the place where you struggle. The treatment plan described by NHLBI usually runs for several weeks and is guided by a provider,

nurse or therapist.

Talk to a healthcare provider if sleep problems are regular, if you repeatedly take more than 30 minutes to fall asleep, wake several times and cannot return to sleep, or feel sleepy during the day. Loud snoring, gasping, choking sounds, pauses in breathing, a strong urge to move tingling legs, or sudden muscle weakness also point beyond an overbusy mind. Sleep studies can measure brain waves, breathing, heart rate, oxygen levels and movement when a disorder is suspected.

For the coming week, choose observation over combat. Give the evening a quieter edge, make the bedroom less stimulating and note what accompanies the difficult nights. If sleep does not improve-or daytime sleepiness is affecting attention, functioning or driving-take that record to a healthcare provider. You do not have to win an argument with your mind before you are allowed to ask for help.

Where this comes from

- Sleep Disorders

  MedlinePlus, U.S. National Library of Medicine · https://medlineplus.gov/sleepdisorders.html
- Insomnia

  MedlinePlus, U.S. National Library of Medicine · https://medlineplus.gov/insomnia.html
- Sleep Disorder Treatments - Sleep Disorder Treatments | NHLBI, NIH

  National Heart, Lung, and Blood Institute · https://www.nhlbi.nih.gov/health/sleep-disorder-treatments
- Sleep Studies - Sleep Studies | NHLBI, NIH

  National Heart, Lung, and Blood Institute · https://www.nhlbi.nih.gov/health/sleep-studies
- About Sleep | CDC

  Centers for Disease Control and Prevention · https://www.cdc.gov/sleep/about/index.html
- A systematic review and meta-analysis of poor sleep, insomnia symptoms and stress in undergraduate students.

  Sleep medicine reviews, 2022 · https://pubmed.ncbi.nlm.nih.gov/34922108/
- Oral magnesium supplementation for insomnia in older adults: a Systematic Review & Meta-Analysis.

  BMC complementary medicine and therapies, 2021 · https://pubmed.ncbi.nlm.nih.gov/33865376/
- Measuring Subjective Sleep Quality: A Review.

  International journal of environmental research and public health, 2021 · https://pubmed.ncbi.nlm.nih.gov/33530453/



- Coffee, caffeine, and sleep: A systematic review of epidemiological studies and randomized controlled trials.

  Sleep medicine reviews, 2018 · https://pubmed.ncbi.nlm.nih.gov/26899133/
- Efficacy and tolerability of pharmacological treatments for insomnia in adults: A systematic review and network meta-analysis.

  Sleep medicine reviews, 2023 · https://pubmed.ncbi.nlm.nih.gov/36701954/