SLEEP & STRESS

Insomnia Has You Awake. Start Here

Before you buy a sleep aid, sort out what is disrupting your nights and choose the first step that fits.

One bad night can become a campaign

You are awake again, doing the bleak arithmetic of how little night remains. By morning, the temptation is to fix everything at once: change your routine, buy a supplement, find a pill, perhaps worry harder. Insomnia encourages urgency. The better first move is slower: work out what kind of problem you are actually having.

Trouble falling asleep is only one version. You may fall asleep easily but wake repeatedly, lie awake for much of the night or surface earlier than you intended. The result can be too little sleep, poor-quality sleep and the feeling that the night never refreshed you. Interrupted sleep matters because sleep quality, not only total hours, supports thinking and daily functioning.

Short-term insomnia often arrives with work stress, family pressure or a traumatic event and lasts days or weeks. Lifestyle changes and good sleep habits often help this kind of disruption. Once the problem lasts a month or longer, it is considered chronic, and the search needs to widen beyond whatever happened yesterday.

That distinction is not a verdict. It is a way to avoid treating every difficult night as the same problem. If this has been brief, begin with the pressures and habits surrounding it. If it has persisted, or keeps returning, it may be time to address the anxiety around sleep and look for another condition, medicine or substance that could be involved.

Look for the disturbance behind the disturbance

Chronic insomnia is often secondary to something else. Medical conditions, pain, depression, anxiety, medicines and other sleep disorders can all disturb sleep. Caffeine, tobacco and alcohol may contribute, as can travel, shift work and an irregular schedule. Sometimes insomnia is the primary problem, and its cause is not well understood.

Stress deserves attention, but not a simplistic explanation. A 2022 review of 34 studies in undergraduate students found moderate associations between stress and both poor sleep quality and insomnia. The studies varied considerably, and most could not establish which came first. Stress and sleeplessness may be entwined without offering you one neat cause to

remove.

Also notice whether the story sounds unlike straightforward insomnia. Loud snoring, gasping, choking sounds or pauses in breathing can point toward sleep apnea. Tingling or prickling in the legs with a strong urge to move them can fit restless legs syndrome. Falling asleep at inappropriate times or experiencing extreme daytime sleepiness may signal another sleep disorder-or simply too little sleep.

This is where the problem stops being only a lifestyle project. Talk with a health care professional if it has lasted a month or longer, regularly disrupts daytime functioning, or comes with breathing pauses, unusual leg sensations or severe daytime sleepiness. Insomnia can impair attention, learning and memory; if you are drowsy while driving, do not treat that as an inconvenience to push through.

Start with the pattern, not the product.

Caffeine is the cleanest experiment

Among the habits worth examining, caffeine has unusually direct evidence behind it. A 2018 systematic review of epidemiological studies and randomized trials found that caffeine typically delayed sleep, reduced total sleep time and sleep efficiency, and worsened people's perception of sleep quality. It also tended to increase wakefulness and arousals.

Timing and amount both mattered in the studies, but sensitivity varied sharply from person to person. Older adults may be more sensitive than younger adults, while substantial differences also appeared among young people. Most research involved men in Western countries, so its findings do not map perfectly onto everyone.

That variability is useful. You do not need to decide whether caffeine is universally "bad." You need to notice whether it belongs in your own chain of events. Coffee and other caffeinated foods and drinks are reasonable places to look before adding a product intended to counteract their effects.

Alcohol belongs on the list of possible contributors too, although the supplied evidence does not establish a precise rule for adjusting it. The same is true of tobacco and irregular schedules. Treat these as clues, not moral failures. Your first task is to identify what may be keeping the problem in motion, not to build a flawless life around sleep.

For persistent insomnia, work on the worry

When insomnia becomes ongoing, the bed can acquire a second burden: anxiety about

whether sleep will happen. Cognitive behavioral therapy, or CBT, can help relieve anxiety linked to chronic insomnia. Relaxation techniques may also reduce anxiety about sleep. The available material here does not specify a do-it-yourself CBT method, so this is one first step to pursue with a qualified professional rather than improvise from a slogan.

A clinician may start with your medical and sleep history and a physical exam, looking for another problem that could be driving the insomnia. A sleep study is sometimes recommended, but not every complaint requires one. Such studies record aspects of sleep and the body's response to sleep problems, including breathing, brain activity, eye movements and heart activity.

Your own sense of sleep still counts. A 2021 review of 49 studies examined several self-report sleep questionnaires and found generally good measurement properties, although some tools had uncertain underlying structures or cutoffs. In other words, a score can organize the conversation, but it does not turn a complicated night into one unquestionable number.

If another condition, medicine or sleep disorder is contributing, addressing that problem matters. This is why persistent insomnia deserves more than an escalating collection of bedtime products. The right first intervention depends on whether you are dealing with stress, a substance, a disrupted schedule, anxiety about sleep or a separate health problem.

A sleep aid is not the obvious beginning

Medicines can help some adults with insomnia, but their benefits and risks differ. A 2023 network meta-analysis of 69 studies involving 17,319 patients compared 20 drugs. Orexin receptor antagonists ranked well across several sleep outcomes, while benzodiazepine-like drugs improved multiple outcomes over placebo but carried a higher safety risk. The long-term adverse effects of all the medications remained unclear.

That is a reason for a risk-and-benefit conversation, not for choosing a drug from a ranking. Most sleep-aid medicines are generally intended for short-term use, and any medicine or supplement should be discussed with your health care professional. The shape of your insomnia-difficulty starting sleep, waking during the night or both-also matters when options are considered.

Melatonin is often treated as the gentler default, but the evidence for adults with chronic insomnia is disappointing. A 2022 review of 24 randomized trials found that, in adults without a related condition, melatonin did not significantly improve time to fall asleep, total sleep time or sleep efficiency. Evidence in adults with another condition was too limited to settle the question.

For the next week, resist turning your nights into a full renovation project. Notice whether caffeine, alcohol, an irregular schedule, stress or a medicine belongs in the picture. If the problem is persistent or daytime life is becoming unsafe, make the next step a clinical conversation. Start with the pattern, not the product.

Where this comes from

- Sleep Disorders

  MedlinePlus, U.S. National Library of Medicine · https://medlineplus.gov/sleepdisorders.html
- Insomnia

  MedlinePlus, U.S. National Library of Medicine · https://medlineplus.gov/insomnia.html
- Healthy Sleep (PDF)

  National Heart, Lung, and Blood Institute · https://www.nhlbi.nih.gov/files/docs/public/sleep/healthy_sleep.pdf
- A systematic review and meta-analysis of poor sleep, insomnia symptoms and stress in undergraduate students.

  Sleep medicine reviews, 2022 · https://pubmed.ncbi.nlm.nih.gov/34922108/
- Measuring Subjective Sleep Quality: A Review.

  International journal of environmental research and public health, 2021 · https://pubmed.ncbi.nlm.nih.gov/33530453/
- Efficacy and tolerability of pharmacological treatments for insomnia in adults: A systematic review and network meta-analysis.

  Sleep medicine reviews, 2023 · https://pubmed.ncbi.nlm.nih.gov/36701954/
- Efficacy of melatonin for chronic insomnia: Systematic reviews and meta-analyses.

  Sleep medicine reviews, 2022 · https://pubmed.ncbi.nlm.nih.gov/36179487/
- Coffee, caffeine, and sleep: A systematic review of epidemiological studies and randomized controlled trials.

  Sleep medicine reviews, 2018 · https://pubmed.ncbi.nlm.nih.gov/26899133/