PROCEDURE

Td (tetanus, diphtheria) vaccine - what you need to know

The Td vaccine is an injection that boosts protection against tetanus and diphtheria.

PURPOSE
Prevent tetanus and diphtheria by boosting immunity

PROCEDURE TYPE
Therapeutic prevention; minimally invasive intramuscular injection

WHERE IT IS DONE
Outpatient clinic, pharmacy, or bedside

ANAESTHESIA
None

HOW LONG IT TAKES
About one to two minutes

PREPARATION
Share vaccine history, allergies, illness, pregnancy, and medicines

RECOVERY
Same day, with no routine driving restriction

MAIN RISKS
Arm pain, fever, fainting, severe allergic reaction

RESULTS
No routine test; protection strengthens over about two weeks

Overview

The Td vaccine protects against tetanus and diphtheria. It is usually given as an injection into a muscle in your upper arm.

Tetanus can cause painful muscle tightening and breathing problems. The bacteria usually enter through broken skin, such as a cut or puncture wound. Tetanus does not spread from person to person.

Diphtheria is a contagious infection that can affect the throat, breathing, heart, and nerves.

The vaccine contains harmless, inactivated forms of toxins made by the bacteria. These are called toxoids . They train your immune system without causing tetanus or diphtheria. Td does not contain live bacteria.

Why the procedure is performed

Td is used to maintain or complete protection against tetanus and diphtheria. You may be offered it:

For wound care, the timing depends on the wound and your vaccine history. A booster may be recommended after five years for a dirty or major wound, or after 10 years for a clean, minor wound.

Td does not protect against pertussis, also called whooping cough. Tdap protects against tetanus, diphtheria, and pertussis.

- As a routine booster, usually every 10 years
- As part of a catch-up series if you were never fully vaccinated
- After certain wounds if your last tetanus-containing vaccine was too long ago
- When the pertussis component of Tdap should not be given

Who should and should not have it

Td may be suitable for people aged 7 years and older who need a tetanus and diphtheria booster or catch-up dose. In many situations, either Td or Tdap can be used after a person has received at least one Tdap dose.

Td may not be the best choice if you have never received Tdap and can safely receive it. Tdap is also specifically recommended during each pregnancy, usually between 27 and 36 weeks, because it helps protect the newborn from pertussis.

Td should not be given if you had a life-threatening allergic reaction after a previous tetanus- or diphtheria-containing vaccine, or if you have a severe allergy to one of its ingredients.

The injection may be delayed if you have a moderate or severe illness. A minor illness, such as a mild cold, is not usually a reason to wait.

Your healthcare team should review the timing carefully if you previously had:

Children younger than 7 generally receive age-specific vaccines such as DTaP rather than Td.

- Severe pain and swelling after a tetanus-containing vaccine
- An Arthus reaction, a severe local immune reaction
- Guillain-Barré syndrome within six weeks of a tetanus-containing vaccine

How to prepare

Follow the treating facility's own instructions, as they take priority over general information.

Fasting is not required. You can usually eat, drink, and take your regular medicines. Do not stop a blood thinner or other prescribed medicine unless your healthcare team tells you to.

Bring your vaccine record if available. Tell the person giving the injection about your allergies, current illness, pregnancy, previous vaccine reactions, bleeding problems, and medicines.

Wear clothing that gives easy access to your upper arm. If you tend to faint around needles, say so before the injection. It can be given while you are seated or lying down.

What happens during the procedure

You will sit or lie in a safe position. The skin over your upper arm may be cleaned.

A healthcare professional inserts a small needle into the muscle and injects the vaccine. The injection itself takes only a few seconds. No anaesthetic or sedation is needed.

You may be asked to remain nearby for a short observation period afterward. This allows staff to help if you feel faint or have an immediate allergic reaction.

Recovery and what to expect afterward

Your arm may feel sore, tender, or heavy. A cool compress and gentle movement may help. Ask your healthcare team which pain medicine is appropriate for you if you need one.

You can eat and drink normally. Staying hydrated may help if you feel light-headed.

Most people can continue ordinary activities, including exercise, as they feel able. Avoid pressing or repeatedly rubbing the injection site.

You can usually drive after Td because it does not involve sedation. Do not drive if you feel faint, dizzy, or unwell.

Most people can return to work or school the same day. Arm discomfort or tiredness may make it more comfortable to reduce strenuous activity briefly.

Risks and possible complications

Common reactions are usually mild and improve within a few days. They can include:

Some people faint during or shortly after an injection. Sitting or lying down can prevent an injury from falling.

A large area of arm swelling can occur, especially after closely spaced tetanus-containing vaccines. Incorrect needle placement can occasionally cause lasting shoulder pain or reduced movement.

A severe allergic reaction is possible after any vaccine but is uncommon. Vaccination sites keep medicines and equipment available to treat this reaction.

No vaccine gives complete protection. Td also cannot cause tetanus or diphtheria.

- Pain, redness, or swelling where the injection was given
- Mild fever, headache, or tiredness
- Nausea or reduced appetite

Results and follow up

There is no routine blood test or immediate result after Td. Your immune system responds over the following days and weeks.

If you are completing a primary vaccine series, you will need additional doses on a planned schedule. Keep a record of the vaccine name and date.

After a high-risk wound, people with an incomplete or uncertain vaccine history may also need tetanus immune globulin. This provides temporary antibodies while the vaccine builds longer-term protection. Wound cleaning and other treatment may still be needed.

Routine follow-up is not usually required unless another vaccine dose has been scheduled or you have a reaction that needs review.

Alternatives

Tdap is the main vaccine alternative. It protects against the same two diseases as Td and also protects against pertussis. Tdap is preferred in certain situations, including during each pregnancy and when an adult has never previously received it.

For many routine boosters, either Td or Tdap may be used. Tdap provides broader protection, but its pertussis component may not be suitable for someone with a specific medical reason to avoid it.

There is no tablet, dietary measure, or natural remedy that replaces vaccination against

tetanus and diphtheria. Wound cleaning reduces contamination but does not provide future immunity. Antibiotics like amoxicillin or doxycycline are not used as a substitute for tetanus vaccination.

Choosing not to be vaccinated avoids expected injection reactions but leaves you with less protection. If you are unsure, ask how your vaccine history, wound type, age, and pregnancy status affect the choice.

When to call your healthcare team

Contact your healthcare team for advice if:

These concerns usually need prompt advice but are not automatically emergencies.

Call emergency services immediately for trouble breathing, swelling of the face or throat, widespread hives, a rapid heartbeat with severe dizziness, collapse, or loss of consciousness. These can be signs of a severe allergic reaction.

- Pain, redness, or swelling is getting worse rather than improving
- A fever is high, persistent, or accompanied by significant illness
- Shoulder pain or reduced movement continues
- You develop unusual weakness, numbness, or tingling
- You are unsure when your next dose is due

Common questions

Will I be asleep for the Td vaccine?

No. Td is a quick injection and does not require sedation or anaesthesia. You remain awake throughout.

Can I drive myself home?

Usually, yes. The vaccine does not impair alertness. Wait until you feel well, and do not drive if you are dizzy or faint.

How painful is the injection?

You may feel a brief pinch or sting. Arm soreness afterward is common and may be more noticeable when you lift the arm.

When will I know it worked?

There is usually no test. Your immune response builds over about two weeks, although full protection may require completion of a vaccine series.

What is the difference between Td and Tdap?

Both protect against tetanus and diphtheria. Tdap also protects against pertussis, or whooping cough. Your vaccine history and reason for vaccination determine which is more appropriate.

Do I need Td after every cut or puncture wound?

No. The need for vaccination depends on the type of wound, when you last received a tetanus-containing vaccine, and whether you completed the primary series. Seek timely advice for deep, contaminated, crushed, burned, or puncture wounds.

Sources

- Centers for Disease Control and Prevention. Td Vaccine: What You Need to Know . Vaccine Information Statement, January 31, 2025.
- Centers for Disease Control and Prevention. Adult Immunization Schedule Notes: Tetanus, Diphtheria, and Pertussis Vaccination . 2025.
- Centers for Disease Control and Prevention. Clinical Guidance for Wound Management to Prevent Tetanus . 2025.
- Havers FP, Moro PL, Hunter P, Hariri S, Bernstein H. Use of Tetanus Toxoid, Reduced Diphtheria Toxoid, and Acellular Pertussis Vaccines: Updated Recommendations of the Advisory Committee on Immunization Practices-United States, 2019 . MMWR Morbidity and Mortality Weekly Report. 2020;69:77-83.
- U.S. Food and Drug Administration. TENIVAC: Tetanus and Diphtheria Toxoids Adsorbed Prescribing Information . Sanofi Pasteur.
- Centers for Disease Control and Prevention. General Best Practice Guidelines for Immunization: Contraindications and Precautions . Advisory Committee on Immunization Practices.