PROCEDURE

Venipuncture

Venipuncture uses a needle to collect blood from a vein for testing or treatment.

PURPOSE
Collecting blood for laboratory testing or therapeutic removal

PROCEDURE TYPE
Diagnostic and therapeutic; minimally invasive needle access

WHERE IT IS DONE
Outpatient clinic, laboratory, blood centre, or bedside

ANAESTHESIA
None; topical numbing cream sometimes used

HOW LONG IT TAKES
1-30 minutes, based on testing or blood removal

PREPARATION
Usually none; some blood tests require fasting

RECOVERY
Same day, with no routine driving restriction

MAIN RISKS
Bruising, bleeding, fainting, and infection

RESULTS
Some within minutes; most laboratory tests take hours to days

Overview

Venipuncture means placing a needle into a vein. It is most often used to collect a blood sample for laboratory testing. It may also be used to remove blood as treatment, which is called therapeutic phlebotomy.

Blood is usually taken from a vein near the inside of your elbow. A vein in your hand may be used if needed. The needle may be connected to collection tubes, a syringe, or a blood collection bag.

A routine blood draw usually takes only a few minutes. Removing a larger amount of blood for treatment takes longer.

Why the procedure is performed

A blood sample can provide information about blood cells, organs, hormones, nutrients, infections, medicines, and many other parts of your health. Venipuncture may be used to:

Therapeutic venipuncture removes a prescribed amount of blood. It may be used for conditions in which the body has too many red blood cells or stores too much iron. The reason, amount removed, and schedule are specific to the condition being treated.

- Help investigate symptoms or monitor a health condition
- Check how well a treatment is working
- Measure medicine levels or possible side effects
- Screen for certain conditions before surgery, pregnancy care, or blood donation
- Collect blood cultures when a bloodstream infection is suspected

Who should and should not have it

Venipuncture is suitable for most people when a blood test or prescribed blood removal is expected to affect care. Before agreeing, you can ask which tests are being ordered, what they may show, and how the results could change your treatment.

There are few absolute reasons to avoid a necessary blood draw. However, a particular vein or arm may not be appropriate. Staff will generally avoid skin that is infected, badly injured, burned, or covered by a large bruise.

Tell the person drawing your blood if you have:

These issues do not always prevent venipuncture. They may affect the site chosen, the equipment used, or how long pressure is applied afterward.

Do not stop a blood thinner like warfarin or apixaban on your own. Most routine blood tests can still be performed, although you may need longer pressure over the site.

Therapeutic blood removal may be delayed if you are unwell, dehydrated, have low blood pressure, or have another condition that makes removing blood unsafe at that time. Your treating team should assess this before proceeding.

- A dialysis fistula or graft in either arm
- An intravenous line or other vascular device
- Lymphoedema, meaning persistent swelling caused by lymph fluid
- Previous lymph node surgery, including surgery related to breast cancer

- A bleeding disorder or a history of prolonged bleeding
- A tendency to faint during needles or blood tests
- An allergy to skin antiseptics, adhesives, or latex

How to prepare

The treating facility's own instructions take priority because preparation depends on the tests being performed.

Most blood tests require no special preparation. For some tests, you may be asked not to eat for a set period. Plain water may still be allowed and can make veins easier to access, but follow the instructions you were given.

Take regular medicines unless your healthcare team has told you otherwise. Ask whether supplements, smoking, caffeine, exercise, or the timing of a medicine could affect your particular test.

Wear clothing that allows easy access above the elbow. Tell staff if you are anxious, have difficult veins, or have fainted before. You may be able to lie down for the procedure. A topical numbing cream can sometimes be used, but it must usually be applied in advance.

What happens during the procedure

Staff will confirm your identity and may ask you to state your name and date of birth. They will check the requested tests and label the collection tubes according to local safety procedures.

You will sit or lie down. A tight band called a tourniquet is placed around your upper arm to make the veins easier to see. You may be asked to close your hand, but repeated forceful fist pumping can affect some test results.

The skin is cleaned with an antiseptic. A sterile needle is then placed into the vein. You may feel a brief sharp pinch or pressure.

Blood flows into one or more tubes. Different tubes contain different substances needed for particular tests. The tourniquet is released, the needle is removed, and pressure is applied with gauze. A small dressing is then placed over the site.

If blood is being removed as treatment, a larger collection bag may be used. Staff may monitor how you feel during and after the procedure.

Recovery and what to expect afterward

Mild soreness or tenderness can occur where the needle entered. A small bruise is also common and should gradually fade. Keeping firm pressure on the site immediately afterward helps limit bleeding and bruising.

You can usually eat and drink normally unless another test or procedure requires restrictions. If you feel light-headed, sit or lie down and have a drink or snack when permitted.

Normal daily activity is usually safe. Avoid heavy lifting with the affected arm for a few hours if the site is sore, bleeding, or bruised. After therapeutic blood removal, you may be advised to rest, drink extra fluids, and avoid strenuous exercise for the remainder of the day.

You can normally drive after a routine blood draw. Do not drive if you feel faint, weak, or unwell, or if you received a sedating medicine for another procedure.

Most people can return to work immediately. After therapeutic blood removal, physically demanding work may need to wait until you feel fully recovered.

Risks and possible complications

Venipuncture is generally a low-risk procedure. Possible problems include:

A sample may occasionally clot, become damaged, or contain too little blood for testing. If this happens, another sample may be needed. This does not necessarily mean there is a problem with your health.

- Pain, bruising, or a collection of blood under the skin
- Continued bleeding, especially with blood thinners or bleeding disorders
- Dizziness, nausea, or fainting
- Difficulty obtaining the sample, requiring another attempt
- Infection at the needle site
- Irritation or injury to a nearby nerve
- Accidental puncture of an artery

Results and follow up

The timing of results depends on the test. Some urgent tests are available within minutes or hours. Many routine results take one or more days. Specialized tests, cultures, or tests sent to

another laboratory may take longer.

Ask when and how you will receive the results. Some facilities release them through an online portal before a clinician has reviewed them. A result outside the laboratory's reference range does not always mean you have a medical problem. Results must be interpreted alongside the reason for testing and other health information.

After therapeutic blood removal, your team may arrange repeat blood counts, iron tests, or other monitoring. Keep routine follow-up appointments even if you feel well.

Alternatives

The alternatives depend on why blood is needed.

A finger-prick or heel-prick sample uses blood from tiny vessels near the skin. It requires less blood and may be easier, but it cannot replace venous blood for many tests. Results may also be more affected by collection technique.

Some health questions can be investigated with urine, saliva, stool, imaging, or physical examination. These options measure different things and may provide less direct or less complete information than a venous blood test.

If you already have a suitable intravenous or central line, blood may sometimes be collected from it. This avoids another needle puncture but can produce misleading results if the sample is contaminated by intravenous fluid or medicine. Handling a central line also carries infection risks.

For selected iron-overload conditions, medicines that bind iron may be an alternative to therapeutic blood removal. They have their own side effects and are not interchangeable for every condition. Erythrocytapheresis selectively removes red blood cells but requires specialist equipment and access.

When to call your healthcare team

Routine follow-up: Contact the team responsible for the test if results do not arrive within the expected time, you are unsure how to access them, or you need help understanding the next step.

Seek prompt medical advice if the needle site develops increasing redness, warmth, swelling, worsening pain, drainage, or a large and painful bruise. Also seek advice for persistent numbness, tingling, weakness, or bleeding that does not stop with continuous firm pressure.

Seek emergency help for heavy uncontrolled bleeding, fainting with incomplete recovery, severe weakness, chest pain, trouble breathing, or swelling of the face, lips, or tongue.

Common questions

Will I be asleep?

No. Routine venipuncture does not require sedation or general anaesthesia. You remain awake. A numbing cream may be an option if needles are particularly difficult for you.

Can I drive home?

Usually, yes. Do not drive if you feel dizzy or faint, or if you received sedation for a separate procedure.

How painful is it?

Most people feel a brief sharp pinch as the needle enters, followed by mild pressure. Tell the person drawing blood if you feel severe, burning, or electric-shock-like pain.

When will I know my results?

Some results are ready within minutes or hours. Routine laboratory results often take one or more days, while cultures and specialized testing may take longer.

What happens if my veins are hard to find?

Staff may warm your arm, lower it, use a smaller needle, or try another site. An experienced practitioner or ultrasound guidance may be requested when access remains difficult.

Can I eat before a blood test?

Usually, but some tests require fasting. Follow the instructions for your specific tests rather than fasting automatically.

Sources

- World Health Organization. WHO Guidelines on Drawing Blood: Best Practices in Phlebotomy . 2010.
- Clinical and Laboratory Standards Institute. GP41: Collection of Diagnostic Venous Blood Specimens . Seventh edition. 2017.

- Simundic AM, Bölenius K, Cadamuro J, and colleagues. Joint EFLM-COLABIOCLI recommendation for venous blood sampling. Clinical Chemistry and Laboratory Medicine . 2018;56(12):2015-2038.
- Centers for Disease Control and Prevention. Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings . 2007, with subsequent updates.
- National Institute for Health and Care Excellence. Infection Prevention and Control: Quality Standard QS61 . Updated 2014.