SYMPTOM

# Voice problems

Voice problems are changes that make your voice sound hoarse, weak, strained, breathy, or different from usual.

**WHAT IT FEELS LIKE**
A hoarse, weak, strained, breathy, or uneven voice

**WHERE IT IS FELT**
Throat and voice box in the neck

**MOST COMMON CAUSES**
Viral laryngitis, voice overuse, throat irritation

**WHEN IT IS URGENT**
Sudden change with swallowing trouble or new neck swelling

**USUALLY LASTS**
Days to weeks, based on irritation or underlying cause

**SEE A DOCTOR IF**
Lasts beyond three to four weeks or keeps returning

## Overview

A voice problem is any change in how your voice sounds or works. It may become rough, breathy, weak, strained, unusually high or low, or difficult to produce. Complete loss of the voice is called aphonia. Other voice changes are often described medically as dysphonia.

Your voice is made when air passes through the vocal folds in your voice box, or larynx. The vocal folds vibrate and produce sound. Swelling, dryness, muscle tension, injury, or changes in how the folds move can affect that sound.

Most short-lived voice problems follow a cold, heavy voice use, or exposure to an irritant. A persistent or unexplained change should be assessed because there are many possible causes.

## What it may feel or look like

Your voice may sound:

Speaking may tire your voice quickly. You might need extra effort to be heard, lose your voice

later in the day, or have difficulty singing notes that were previously comfortable.

Other symptoms can include a dry or sore throat, coughing, frequent throat-clearing, a feeling of a lump in the throat, or discomfort when speaking. Trouble forming words, slurred speech, and difficulty understanding language are not simply voice problems and may need separate assessment.

- Hoarse, raspy, or rough
- Breathy, faint, or weak
- Tight, strained, or effortful
- Lower or higher than usual
- Unsteady, shaky, or broken

## Common causes

Common causes include viral laryngitis, which is inflammation of the voice box during or after a cold. Voice overuse from shouting, singing, teaching, or speaking for long periods can also irritate the vocal folds.

Smoke, vaping, dry air, dust, chemical fumes, and allergies may irritate the throat. Acid reflux can contribute in some people, although hoarseness alone does not prove that reflux is the cause. Muscle tension around the voice box may make the voice feel strained, especially during stress or prolonged voice use.

Less common causes include small, noncancerous growths such as vocal fold nodules, polyps, or cysts. These may develop after repeated strain or an injury to a vocal fold. Age-related thinning of the vocal folds can make the voice weaker or breathier.

Some medicines can dry or irritate the throat. Inhaled medicines may occasionally affect the voice. Thyroid problems, nerve conditions, and reduced movement or paralysis of a vocal fold are other possibilities. A nerve-related change can sometimes follow neck or chest surgery, a breathing tube, or an injury.

Causes needing prompt assessment include bleeding into a vocal fold after forceful shouting or singing, a serious infection, or a growth affecting the voice box. Cancer is not the usual cause of hoarseness, but persistent changes are more concerning when accompanied by a neck lump, coughing blood, unexplained weight loss, or a history of smoking.

## When to seek medical care

Call emergency services if the voice change occurs with:

Seek same-day medical advice for a sudden voice change after a neck injury, surgery, or placement of a breathing tube, particularly if you also have choking or trouble swallowing. Sudden voice loss after forceful singing, shouting, or coughing also deserves prompt assessment.

Book a routine appointment if the change lasts longer than three to four weeks, keeps returning, or interferes with work or daily life. Arrange an earlier review if you have pain when speaking, a neck lump, coughing blood, ongoing swallowing difficulty, unexplained weight loss, or a weakened immune system.

Children should be assessed promptly if a voice change comes with noisy breathing, drooling, difficulty swallowing, or unusual sleepiness.

- Difficult or noisy breathing
- Inability to swallow saliva
- Rapid swelling of the tongue, throat, or neck
- Facial drooping, one-sided weakness, confusion, or sudden speech difficulty

## How the cause is found

A healthcare professional will ask when the change began, whether it followed an illness or heavy voice use, and whether you smoke, vape, take medicines, or have reflux or swallowing symptoms. Mention recent surgery, neck injury, or breathing-tube placement.

They may examine your mouth, throat, neck, breathing, and speech. If the problem persists or there are concerning features, you may be referred to an ear, nose, and throat specialist.

The vocal folds can be viewed using laryngoscopy. This usually involves passing a thin, flexible camera through the nose after numbing spray is applied. Imaging scans are not usually the first test for an isolated voice change; the vocal folds are generally examined first.

## Treatment and self-care

Treatment depends on the cause. For a short-lived illness or mild irritation, use your voice gently and avoid shouting, singing, or long conversations. Whispering is not ideal because it can still strain the voice. Speak quietly at a comfortable pitch instead.

Drink enough fluids and consider a clean humidifier if the air is dry. Avoid smoking, vaping, and

smoky environments. Try not to clear your throat repeatedly; sipping water or swallowing may be less irritating.

Antibiotics like amoxicillin or doxycycline do not help viral laryngitis. Steroids and reflux medicines are not routine treatments for every voice problem and should not be started solely because the voice is hoarse without appropriate assessment.

Persistent problems may be treated with voice therapy from a speech and language therapist. Treatment for a vocal fold growth, paralysis, or another structural problem may involve a procedure or surgery. Any medicine thought to be contributing should be reviewed rather than stopped without advice.

## What to expect

Voice changes from a cold or brief overuse often improve within one or two weeks. Recovery may take longer if you keep straining the voice or remain exposed to an irritant.

A persistent problem does not necessarily mean something serious, but it is less likely to resolve without identifying the cause. Voice therapy and changes in voice habits can help many long-term problems. Some conditions need ongoing specialist care or a procedure.

## Common questions

Should I whisper to rest my voice?

Usually not. Whispering can create tension around the voice box. Use less voice overall and speak gently at a comfortable volume.

Can reflux cause hoarseness without heartburn?

It can, but hoarseness, throat-clearing, and a lump sensation have several possible causes. These symptoms alone cannot confirm reflux.

Do warm drinks help?

Warm, non-irritating drinks may soothe dryness or discomfort, but they do not directly reach the vocal folds. Avoid drinks that seem to worsen reflux or irritation.

Could stress affect my voice?

Yes. Stress can increase muscle tension and alter breathing, making the voice sound tight or weak. A clinician should still assess persistent changes rather than assuming stress is the only cause.

## Sources

- Stachler RJ, Francis DO, Schwartz SR, et al. Clinical Practice Guideline: Hoarseness (Dysphonia), 2018 update . Otolaryngology-Head and Neck Surgery . 2018;158(1 Suppl):S1-S42.
- National Institute on Deafness and Other Communication Disorders. Hoarseness .
- National Institute on Deafness and Other Communication Disorders. Taking Care of Your Voice .
- American Speech-Language-Hearing Association. Voice Disorders Practice Portal .
- NHS. Laryngitis .