TEST

HCG blood test - quantitative

A quantitative HCG blood test measures the amount of pregnancy-related HCG hormone in your blood.

WHAT IT MEASURES
Amount of human chorionic gonadotropin hormone in blood

SAMPLE
Blood

FASTING NEEDED?
No

HOW LONG IT TAKES
5 to 10 minutes

RESULTS IN
Same day to two days depending on laboratory

USED FOR
Confirming and monitoring pregnancy or HCG-producing conditions

MAIN LIMITATION
One result cannot show pregnancy location or healthy development

Overview

A quantitative HCG blood test measures the amount of human chorionic gonadotropin, or HCG, in your blood. It gives a number rather than a simple positive or negative result.

HCG is made mainly by tissue that develops into the placenta during pregnancy. It usually becomes detectable in blood after implantation, when a fertilized egg attaches to the lining of the uterus.

Small amounts can sometimes come from other sources. These include the pituitary gland, particularly around or after menopause, and certain uncommon tumors. The result must therefore be interpreted in context.

A quantitative blood test can detect lower HCG amounts than many urine pregnancy tests. However, one measurement cannot confirm where a pregnancy is located, whether it is developing normally, or exactly how long you have been pregnant.

Why the test is ordered

The test is often ordered to confirm a possible pregnancy when an early or more precise result is needed. It may also help assess pregnancy-related symptoms or follow changes over time.

Reasons for testing include:

An ectopic pregnancy develops outside the uterus, most often in a fallopian tube. HCG measurements can support the assessment, but ultrasound and your symptoms are also important.

- Checking for pregnancy before some procedures or treatments
- Investigating pain or bleeding during early pregnancy
- Monitoring a pregnancy when miscarriage or ectopic pregnancy is possible
- Following HCG after a miscarriage, abortion, delivery, or ectopic pregnancy
- Monitoring fertility treatment that involves pregnancy testing
- Assessing gestational trophoblastic disease, which involves abnormal placental tissue
- Monitoring certain ovarian, testicular, or other HCG-producing tumors

Who should have the test

You may be offered this test if a urine test is unclear, if testing is being done very early, or if your care team needs an actual HCG level rather than a positive or negative answer.

It may also be appropriate if you have early-pregnancy pain or bleeding, have had fertility treatment, or need follow-up after a pregnancy ends. People being assessed or monitored for an HCG-producing tumor may have the same test for a different reason.

Quantitative HCG testing is not usually needed to confirm every uncomplicated pregnancy. A home or laboratory urine test may provide enough information in many situations.

How to prepare

You do not need to fast. You can usually eat, drink water, and have coffee as normal.

Tell the person ordering the test about:

Do not stop prescribed medicines or supplements unless you have been told to do so. If repeated tests are planned, using the same laboratory can make the results easier to compare.

- Fertility medicines or injections containing HCG
- Prescription medicines, supplements, and high-dose biotin
- A recent pregnancy, miscarriage, abortion, or delivery
- Kidney disease
- Your last menstrual period, if pregnancy is possible
- Any recent HCG results from another laboratory

What happens during the test

A healthcare professional places a small needle into a vein, usually in your arm. Blood is collected into a tube and sent to a laboratory.

The blood draw usually takes 5 to 10 minutes. The appointment may take longer if you need registration or other tests.

If the result is being used to assess an early pregnancy, another sample may be taken after about 48 hours or at another interval chosen for your situation. The pattern over time often provides more information than one value.

Understanding your results

Results are commonly reported as milli-international units per millilitre, written as mIU/mL, or international units per litre, written as IU/L. These units are numerically equivalent.

Laboratories set their own reference information. Interpretation also varies with age, sex, pregnancy status, timing, medicines, the testing method, and the clinical situation.

A detectable or increasing result can support pregnancy, but it does not prove that the pregnancy is inside the uterus or developing normally. Healthy pregnancies can have different HCG levels and rates of change.

A falling level may occur after a miscarriage, abortion, delivery, or treatment for ectopic pregnancy. Follow-up may continue until the level reaches the target set for your circumstances.

A higher result does not by itself mean that you are carrying twins. Pregnancy dating differences, multiple pregnancy, abnormal placental tissue, and other factors can overlap.

When HCG is used as a tumor marker, results are interpreted with imaging, examination, other blood tests, and sometimes tissue testing. Pregnancy must be considered whenever it is

biologically possible.

What affects the results

Testing too early is the most common reason for a negative result when pregnancy is present. Implantation timing differs, so HCG may not yet have reached the test's detection limit.

Other influences include:

Ordinary food, coffee, time of day, and light daily activity do not usually change serum HCG meaningfully. A recent blood transfusion is not a common cause of an altered result.

- HCG fertility injections: These can cause a positive result until the injected hormone clears
- Recent pregnancy: HCG may remain detectable for a time after delivery, miscarriage, abortion, or ectopic pregnancy treatment
- Different laboratories: Testing systems may measure forms of HCG differently, so results may not match exactly
- Biotin supplements: High doses can interfere with some laboratory methods; the direction of error depends on the system
- Antibody interference: Naturally occurring antibodies can interact with the test and produce a false result
- Very high HCG: Rarely, an extreme concentration overwhelms an assay and produces a falsely low result
- Age and menopause: The pituitary gland may produce small amounts of HCG around or after menopause
- Kidney impairment: Reduced clearance can contribute to persistent low-level results
- HCG-producing conditions: Some tumors and abnormal placental tissue raise HCG without a typical pregnancy

Accuracy and limitations

A quantitative blood test is good at detecting small amounts of HCG, but no test is perfect. Testing before enough HCG has developed can miss a pregnancy. Rare laboratory interference can report HCG when clinically meaningful HCG is not present.

If a result does not fit your symptoms or other findings, the laboratory may repeat it using another testing method. Comparing a blood result with a urine test can sometimes help

identify antibody interference because these antibodies usually do not pass into urine.

One HCG number cannot reliably date a pregnancy. It also cannot show whether the pregnancy is in the uterus, has a heartbeat, or will continue normally.

Repeated measurements can show a trend, but there is overlap between healthy pregnancies, miscarriages, and ectopic pregnancies. Ultrasound is usually the better test for locating a pregnancy and assessing development once it is expected to be visible.

Risks and complications

The main risks are brief discomfort, a small bruise, or minor bleeding where the needle entered. Some people feel faint during or after a blood draw.

What happens next

A routine positive result may be followed by standard pregnancy care. If the pregnancy is very early or the result is uncertain, repeat HCG testing or an ultrasound may be arranged.

For monitoring after a pregnancy ends or during treatment, testing may continue until the expected trend or target is reached. Tumor-marker results are followed according to the condition being assessed and are not interpreted as pregnancy results.

Seek urgent medical assessment if pregnancy is possible and you develop worsening one-sided abdominal or pelvic pain, shoulder-tip pain, fainting, marked weakness, or heavy bleeding. These symptoms need prompt evaluation regardless of the HCG number because they can occur with internal bleeding or ectopic pregnancy.

Mild symptoms, an unexpected test result, or questions about repeat testing can usually be discussed through routine follow-up.

Common questions

Can I drink coffee before the test?

Yes. Fasting is not required, and coffee does not usually affect the result.

Does the time of day matter?

Not usually for a blood HCG test. If levels are being followed, your care team may still arrange samples at consistent intervals.

Can I exercise beforehand?

Normal exercise is unlikely to change the result. If exercise makes you dehydrated or light-headed during blood tests, drink water and rest beforehand.

How soon can a blood test detect pregnancy?

Blood testing can detect pregnancy earlier than many urine tests, but testing too soon can still be negative. Implantation timing and the laboratory's detection limit both matter.

Why do I need another test in about 48 hours?

A second test shows whether HCG is rising, falling, or staying similar. The trend is more informative than one result, although it still cannot locate the pregnancy.

Can the test be positive when I am not pregnant?

Yes, although this is uncommon. Recent pregnancy, HCG fertility medicine, pituitary HCG, laboratory interference, and certain HCG-producing conditions can cause a detectable result.

Sources

- MedlinePlus Medical Encyclopedia: HCG blood test - quantitative
- American College of Obstetricians and Gynecologists: Practice Bulletin No. 193, Tubal Ectopic Pregnancy
- American College of Obstetricians and Gynecologists: Practice Bulletin No. 200, Early Pregnancy Loss
- American College of Obstetricians and Gynecologists: Avoiding Inappropriate Clinical Decisions Based on False-Positive Human Chorionic Gonadotropin Test Results
- Mayo Clinic Laboratories: Chorionic Gonadotropin, Total, Beta-Subunit, Quantitative, Serum
- US Food and Drug Administration: Biotin Interference with Certain In Vitro Diagnostic Tests